

A FAUNISTIC CATALOGUE OF LONGHORNED BEETLES OF MARMARA REGION OF TURKEY (COLEOPTERA: CERAMBYCOIDEA: CERAMBYCIDAE)

| Hüseynin Özdikmen* |

* Department of Biology, Faculty of Science, Gazi University, 06500 Ankara, TURKEY. E-mail: ozdikmen@gazi.edu.tr, ORCID ID: 0000-0001-9568-0093

[Özdikmen, H. 2026. A faunistic catalogue of longhorned beetles of Marmara region of Turkey (Coleoptera: Cerambycoidea: Cerambycidae). Munis Entomology & Zoology, 21 (1): 887-970]

ABSTRACT: This faunistic catalogue presents an updated list of all species-group taxa of longhorned beetles in Marmara region of Turkey with provincial distributions. As a result of this, a total of 332 species-group taxa of 93 subgenera of 98 genera belonging to 51 tribes of 11 subfamilies in one family as Cerambycidae are determined and presented.

KEY WORDS: Cerambycoidea, Cerambycidae, Marmara region, Turkey

Turkey is located in the northern hemisphere between the 36°-42° northern parallel and the 26°-45° eastern meridian on an area of 783 562 km². It is situated between Europe and Asia continents, and creating a link between them. The Asian part is called as Anatolia (or Asia Minor), while the European part is known Thrace or East Thrace. They are separated by the Sea of Marmara, the Bosphorus, and the Dardanelles. Turkey is a peninsula which is encircled by seas on three sides: the Black Sea in the North, the Aegean Sea in the West, and the Mediterranean Sea in the South (Fig. 1).

Turkey is generally divided into seven geographical regions (Fig. 1). Marmara region, and its parts and provinces can be presented as below.

1. Marmara Region (North-West Turkey)

Section A. European Turkey (Thrace or East Threce)

1. Ergene Part (including Edirne and Tekirdağ provinces)
2. Istranca Part (including Kırklareli province)
3. Çatalca Part (including İstanbul province, and Çanakkale province partly)

Section B. Asian Turkey (North-West Anatolia)

1. Kocaeli Part (including Kocaeli and Sakarya provinces, and İstanbul province partly)
2. South Marmara Part (including Bilecik, Yalova, Bursa, Balıkesir and Çanakkale provinces)

The Marmara region is a geographical region of Turkey in North-Western part of Turkey. It is bordered by the Greece to the West, Bulgaria to the North-West, Aegean Sea to the South-West, the Black Sea to the North, the Black Sea region to the East, the Aegean region to the South, and the Central Anatolian region to the Sout-East. The Marmara region has an area of 72 666 km² and is the smallest

region of Turkey. The Marmara region has a humid subtropical climate bordering the Mediterranean climate along the Aegean and Southern Marmara coasts, while the Black Sea coast has an oceanic climate. Summers are warm to hot and moderately dry, while winters are cool, rainy, and sometimes snowy. The coastal climate maintains relatively mild temperatures.

The area includes the whole territories of the Turkish provinces Balıkesir (14583 km²), Bilecik (4179 km²), Bursa (10813 km²), Çanakkale (9817 km²), Edirne (6145 km²), İstanbul (5461 km²), Kırklareli (6459 km²), Kocaeli (3397 km²), Sakarya (4824 km²), Tekirdağ (6190 km²), and Yalova (798 km²). Among them, Çanakkale and İstanbul provinces have a European (Thrace) part and an Asian (Anatolia) part (Fig. 2).

First comprehensive study of longhorned beetles fauna of Marmara region of Turkey was realized by Özdikmen (2008). According to Özdikmen (2008), the longhorned beetles fauna of Marmara region of Turkey was consisted of 264 species of 102 genera of 7 subfamilies.



Figure 1. 81 provinces and 7 regions in Turkey [(1) Marmara region, (2) Black Sea region, (3) Aegean region, (4) Central Anatolian region, (5) Eastern Anatolian region, (6) Mediterranean region, (7) South-Eastern Anatolian region].





Figure 2. Marmara region and its provinces.

With the present updation, all species-group taxa of longhorned beetles in Marmara region of Turkey with their provincial distribution in Marmara region of Turkey are determined and presented below. These species-group taxa are determined on the base of the study of Özdikmen (2008, 2021, 2024, 2025a). This is an updated list with some newly determined taxa.

All higher taxa (superfamily to tribe) are arranged phylogenetical order. Also, all taxa are arranged completely alphabetical order within each tribe an except for synonyms that are given for genus-group and species-group names and ordered to priority.

The currently valid names of the family-group taxa include the name of the author and the year of the publication. Their synonyms are not listed.

The names of the genus-group taxa are given with the name of the author, and the year and page of publication. The type species of all genus-group names are given in their original combination. If the type species is currently regarded as a junior synonym, the valid senior synonym is given in brackets in its original combination.

The names of the species-group taxa are given with the name of the author, and the year and page of publication. For species-group taxa subsequently transferred to another genus, the name of the original genus is given in parentheses, following the page of publication.

The following symbols, all given in square brackets following the page of publication, or the original combination when applicable, are used for taxonomic information: HN for homonyms, RN for replacement names, NO for nomina oblita, NP for nomina protecta, DA for doubtful assignment.

The valid taxa originally described from Turkey are marked with the sign (#) in red color at the end of sentences containing taxa names.

Endemic taxa for Turkey are marked with the sign (***) in red color at the end of sentences containing taxa names.

References were given to primary sources of the family-group, genus-group and species-group names included in the studies of Özdikmen (2021, 2024 and 2025a). Additional new references of Cerambycoidea taxa of Turkey are given at the end of present study.

AN UPDATED FAUNISTIC LIST OF CERAMBYCOIDEA TAXA OF MARMARA REGION OF TURKEY (COLEOPTERA: CERAMBYCIDAE)

Superfamily CERAMBYCOIDEA

Family CERAMBYCIDAE Latreille, 1802

Subfamily PRIONINAE Latreille, 1802

Tribe Ergatini Fairmaire, 1864

Genus **Ergates** Audinet-Serville, 1832: 143

type species *Prionus serrarius* Panzer, 1793 (= *Cerambyx faber* Linnaeus, 1760)

Caratambyx Gistel, 1856: 375 type species *Cerambyx faber* Linnaeus, 1760

faber faber Linnaeus, 1760: 187 (*Cerambyx*)

portitor Schrank, 1781b: 312 (*Cerambyx*)

bulzanensis Laicharting, 1784: 1 (*Prionus*)

serrarius Panzer, 1793: 6 (*Prionus*)

obscurus Olivier, 1795: 26 (*Prionus*)

crenatus Fabricius, 1802: 264 (*Prionus*)

grandiceps Tournier, 1872: 257

alkani Demelt, 1968b: 28

Kocaeli, Bursa, Yalova, Edirne

Tribe Macrotomini J. Thomson, 1861

Subtribe Macrotomina J. Thomson, 1861

Genus **Prinobius** Mulsant, 1842b: 207

type species *Prinobius myardi* Mulsant, 1842b

Prionobius Chevrolat, 1854: 482 [unnecessary emendation]

myardi slamorum Danilevsky, 2012b: 698 [RN]

scutellaris Germar, 1817: 219 [HN] (*Prionus*)

İstanbul (Anatolia), Çanakkale
(Anatolia), İstanbul (Thrace),
Çanakkale (Gelibolu)

Subtribe **Remphanina** Lacordaire, 1868

Genus **Rhaesus** Motschulsky, 1875: 153 [RN]

type species *Rhaesus persicus* Motschulsky, 1875 (= *Prionus serricollis* Motschulsky, 1838)

Rhesus J. Thomson, 1861: 291 [HN] type species *Prionus serricollis* Motschulsky, 1838

serricollis Motschulsky, 1838: 187 (*Prionus*)

robustus C. H. G. Heyden, 1844: 15 (*Aulacopus*)

persicus Motschulsky, 1875: 154

Kocaeli, İstanbul (Anatolia),
Bilecik, Edirne

Tribe **Aegosomatini** J. Thomson, 1861

Genus **Aegosoma** Audinet-Serville, 1832: 162

type species *Cerambyx scabricornis* Scopoli, 1763

Subgenus **Aegosoma** Audinet-Serville, 1832: 162

type species *Cerambyx scabricornis* Scopoli, 1763

scabricorne Scopoli, 1763: 54 (*Cerambyx*)

İstanbul (Anatolia), Balıkesir,
Bilecik, Kırklareli, İstanbul
(Thrace)

Tribe **Prionini** Latreille, 1802

Genus **Mesoprionus** Jakovlev, 1887: 323

type species *Prionus asiaticus* Faldermann, 1837

Subgenus **Trispinicollis** Özdikmen, 2025b: 3001

type species *Prionus besikanus* Fairmaire, 1855

besikanus Fairmaire, 1855: 318 (*Prionus*)

batelkai Sláma, 1996: 75 (*Prionus*)

İstanbul (Anatolia), Bilecik,
Bursa, Çanakkale (Anatolia),
Edirne, İstanbul (Thrace)

Genus **Prionus** Geoffroy, 1762: 198

type species *Cerambyx coriarius* Linnaeus, 1758

Prionellus Casey, 1924: 209 type species *Prionus pocularis* Dalman, 1817

coriarius Linnaeus, 1758: 389 (*Cerambyx*)

tridentatus Linnaeus, 1758: 354 (*Scarabaeus*)

prionus DeGeer, 1775: 59 (*Cerambyx*)

vicinus Jakovlev, 1887: 325

schaufussi Jakovlev, 1887: 326

burdajewiczi Bodemeyer, 1927: 74

Kocaeli, Sakarya, İstanbul
(Anatolia), Balıkesir,
Bilecik, Çanakkale
(Anatolia), Kırklareli,
İstanbul (Thrace),
Çanakkale (Gelibolu)

Subfamily **LEPTURINAE** Latreille, 1802

Tribe **Xylosteini** Reitter, 1913

Genus **Xylosteus** Frivaldszky von Frivald, 1837: 180
type species *Xylosteus spinolae* Frivaldszky von Frivald, 1837

spinolae Frivaldszky von Frivald, 1837: 180
rufiventris Germar, 1844: 16 (*Rhagium*)
merkli Pic, 1910c: 66

Kırklareli

Tribe **Rhamnusiini** Sama, 2009

Genus **Rhamnusium** Latreille, 1829: 130
type species *Callidium salicis* Fabricius, 1787 (= *Cerambyx bicolor* Schrank, 1781a)

bicolor graecum Schaefuss, 1862: 311
limbatum Pic, 1897c: 30

İstanbul (Thrace)

bicolor juglandis Fairmaire, 1866: 276 # ***
anatolicum Pic, 1901b: 10
delagrangei Pic, 1901b: 10
geniculatum Pic, 1901b: 10
obscuripes Pic, 1903d: 163
rufotibiale Pic, 1917a: 3

İstanbul (Anatolia), Bursa,
Çanakkale (Anatolia),
Çanakkale (Gelibolu)

Tribe **Evodinini** Zamoroka, 2022

Genus **Brachyta** Fairmaire, 1865: 185
type species *Leptura interrogationis* Linnaeus, 1758

Subgenus **Fasciobrachyta** Danilevsky, 2014b: 113
type species: *Leptura bifasciata* Olivier, 1800

balcanica Hampe, 1871: 336

Kırklareli

Tribe **Acmaeopsini** Della Beffa, 1915

Genus **Cortodera** Mulsant, 1863: 572
type species *Grammoptera spinosula* Mulsant, 1839: 290 (= *Leptura humeralis* Schaller, 1783)

Leptacmaeops Casey, 1913: 219 type species *Leptura longicornis* Kirby, 1837
Acmaeopsilla Casey, 1913: 240 type species *Acmaeops falsus* LeConte, 1860
Taiwanocarilia Hayashi, 1983: 30 type species *Gaurotes ater* Tamanuki, 1943

differens magdae Danilevsky, 2012a: 916

Kırklareli

flavimana flavimana Walzl, 1838: 471 (*Leptura*) # Kocaeli, İstanbul (Anatolia),
Tekirdağ, Kırklareli,
İstanbul (Thrace)

flavimana inonuensis Danilevsky, 2015c: 1045 # *** Bursa

humeralis humeralis Schaller, 1783: 297 (*Leptura*) Kırklareli
quadriguttata Herbst, 1786: 171 (*Leptura*)
suturalis Fabricius, 1787: 159 (*Leptura*)
quadrinotata Gmelin, 1790: 1873 (*Leptura*)
schalleri Gmelin, 1790: 1874 (*Leptura*)
spinosula Mulsant, 1839: 290 (*Grammoptera*)
inhumeralis Pic, 1892h: 140
discoidalis Pic, 1931b: 6
apicenotata Schmidt, 1951: 12

pumila meltemae Özdkimen, Mercan & Cihan, 2012d: 746 # *** Bilecik, Kırklareli

rufipes rufipes Kraatz, 1876: 344 (*Grammoptera*) # *** Bursa

Genus *Dinoptera* Mulsant, 1863: 494
type species *Leptura collaris* Linnaeus, 1758

collaris Linnaeus, 1758: 398 (*Leptura*) Kocaeli, İstanbul (Anatolia),
Bursa, Kırklareli
ruficollis DeGeer, 1775: 143 (*Leptura*)
sylvestris Geoffroy, 1785: 88 (*Stenocorus*)
morio Fabricius, 1793: 349 (*Leptura*)
carneola Schrank, 1798: 696 (*Leptura*)

Tribe **Pidoniini** Zamoroka, 2022

Genus *Fallacia* Mulsant & Rey, 1863: 180
type species *Fallacia longicollis* Mulsant & Rey, 1863 (= *Grammoptera elegans* Faldermann, 1837)

elegans Faldermann, 1837: 319 (*Grammoptera*) Kırklareli
longicollis Mulsant & Rey, 1863: 180
mingrelica Kraatz, 1879b: 279
lederi Kraatz, 1879b: 280
inapicalis Pic, 1901j: 16
subnotata Pic, 1901j: 16

Tribe **Rhagiini** Kirby, 1837

Genus *Hagrium* Villiers, 1978: 85
type species *Rhagium bifasciatum* Fabricius, 1775

bifasciatum Fabricius, 1775: 183 (*Rhagium*) Kocaeli, Sakarya, İstanbul
ornatum Fabricius, 1775: 183 (*Rhagium*) (Anatolia), Kırklareli
maculatum Goeze, 1777: 462 (*Cerambyx*)

parisinum Geoffroy, 1785: 85 (*Stenocorus*)
anglicum Gmelin, 1790: 1844 (*Cerambyx*)
bicolor Olivier, 1800: 16 (*Stenocorus*)
nigrolineatum Donovan, 1801: 73 (*Leptura*)
bimaculatum Marsham, 1802: 343 (*Leptura*)
ecoffeti Mulsant, 1839: 222 (*Rhagium*)
unifasciatum Mulsant, 1839: 222 (*Rhagium*)
lituratum Fügner, 1891: 201 (*Stenocorus*)
latefasciatum Pic, 1891c: 2 (*Rhagium*)
infasciatum Pic, 1898a: 3 (*Rhagium*)
gravei Hubenthal, 1902: 291 (*Rhagium*)
deyrollei Pic, 1909: 123 (*Rhagium*)
fasciatum Pic, 1910d: 19 (*Rhagium*)
rufum Pic, 1910d: 19 (*Rhagium*)
medionotatum Pic, 1910d: 20 (*Rhagium*)
bistrinotatum Pic, 1914a: 3 (*Rhagium*)

Genus **Megarhagium** Reitter, 1913a: 6
type species *Cerambyx sycophanta* Schrank, 1781a

fasciculatum Faldermann, 1837: 304 (*Rhagium*)
rufipes Motschulsky, 1838: 184 (*Rhagium*)

Kocaeli

mordax DeGeer, 1775: 124 (*Leptura*)
bifasciatum Schrank, 1781a: 137 (*Leptura*) [HN]
linnei Laicharting, 1784: 122 (*Rhagium*)
vulgare Samouelle, 1819: 210 (*Rhagium*)
klenkai Pic, 1915b: 4 (*Rhagium*)
subdilatatum Pic, 1917a: 3 (*Rhagium*)
morvandicum Pic, 1927b: 10 (*Rhagium*)
altajense Podány, 1964b: 15 (*Rhagium*)

Edirne

sycophanta Schrank, 1781a: 137 (*Cerambyx*)
scrutator Olivier, 1800: 10 (*Stenocorus*)
cephalotes Mulsant, 1839: 224 (*Rhagium*)
grandiceps C. G. Thomson, 1866: 50 (*Rhagium*)

Kırklareli

Genus **Rhagium** Fabricius, 1775: 182
type species *Cerambyx inquisitor* Linnaeus, 1758
Hargium Leach, 1819: 210 type species *Cerambyx inquisitor* Linnaeus, 1758
Allorhagium Kolbe, 1884: 270 type species *Cerambyx inquisitor* Linnaeus, 1758

inquisitor inquisitor Linnaeus, 1758: 393 (*Cerambyx*)
nubecula Bergsträsser, 1778: 26 (*Cerambyx*)
indagator Fabricius, 1787: 145
minutum Fabricius, 1787: 146
exile Gmelin, 1790: 1844 (*Cerambyx*)
investigator Mulsant, 1839: 227
iberonis Ericson, 1916: 240

Sakarya, İstanbul (Anatolia),
Balıkesir, Bursa, Edirne,
İstanbul (Thrace)

Genus **Stenocorus** Geoffroy, 1762: 221
type species *Leptura meridiana* Linnaeus, 1758

Subgenus **Anisorus** Mulsant, 1862: 467
type species *Cerambyx quercus* Götze, 1783

quercus quercus Götze, 1783: 74 (*Cerambyx*)
humeralis Fabricius, 1787: 158 (*Leptura*) [HN]
dispar Panzer, 1794: 2
bajus Schrank, 1798: 692 (*Leptura*)
discolor Fleischer, 1887a: 237 (*Toxotus*)
unicolor Fleischer, 1887b: 272 (*Toxotus*) [unnecessary RN]
magdalenae Pic, 1891c: 4 (*Toxotus*)
punctipennis Reitter, 1914: 178 (*Stenochorus*)

Edirne

Subgenus **Stenocorus** Geoffroy, 1762: 221
type species *Leptura meridiana* Linnaeus, 1758
Endemus Gistel, 1848b: xi [unnecessary RN]
Minaderus Mulsant, 1862: 467 type species *Leptura meridiana* Linnaeus, 1758
Toxotus Dejean, 1821: 112 type species *Leptura meridiana* Linnaeus, 1758

meridianus Linnaeus, 1758: 398 (*Leptura*)
grandis Poda von Neuhaus, 1761: 38 (*Leptura*)
ruficrus Scopoli, 1763: 48 (*Leptura*)
chrysogaster Schrank, 1781a: 132 (*Cerambyx*)
cantharinus Herbst, 1784: 93 (*Rhagium*)
splendens Laicharting, 1784: 137 (*Leptura*)
geniculatus Geoffroy, 1785: 86
sericeus Olivier, 1800: 20
laevis Olivier, 1800: 21
rufiventris Marsham, 1802: 341 (*Leptura*)
glabratus Latreille, 1804: 311 (*Leptura*)
luteolus L. Petagna, 1819: 37
lacordairii Pascoe, 1867: lxxxiv (*Toxotus*)
bilineatus Pic, 1908a: 58 (*Stenochorus*)
brevelineatus Pic, 1941a: 5
brevesignatus Pic, 1941a: 5
postsignatus Pic, 1941d: 13

Edirne

Tribe **Lepturini** Latreille, 1802

Genus **Allosterna** Mulsant, 1863: 576
type species *Leptura tabacicolor* DeGeer, 1775
Allosterna Stierlin, 1898: 479 [unjustified emendation]

tabacicolor tabacicolor DeGeer, 1775: 139 (*Leptura*)
chrysomeloides Schrank, 1781a: 158 (*Leptura*)
solstitialis Herbst, 1784: 103 (*Leptura*)
genii Gmelin, 1790: 1877 (*Leptura*)
laevis Fabricius, 1793: 340 (*Leptura*)
dispar Pic, 1892h: 140
bicoloripes Pic, 1914b: 65 (*Allosterna*)

İstanbul (Anatolia), Balıkesir,
Kırklareli, İstanbul (Thrace)

Genus **Anastrangalia** Casey, 1924: 280
type species *Leptura sanguinea* LeConte, 1859
Marthaleptura K. Ohbayashi, 1963: 9 type species *Leptura scotodes* Bates, 1873

- dubia dubia*** Scopoli, 1763: 47 (*Leptura*) Bilecik, Bursa
limbata Laicharting, 1784: 166 (*Leptura*)
notata Olivier, 1800: 11 (*Leptura*)
cincta Fabricius, 1801: 356 (*Leptura*)
chamomillae Fabricius, 1801: 359 (*Leptura*)
luctuosa Mulsant, 1839: 278 (*Leptura*)
basinotata Pic, 1932: 31 (*Leptura*)
graeca Pic, 1932: 31 (*Leptura*)
birubronotata Pic, 1941b: 1 (*Leptura*)
planeti Pic, 1945a: 5
curierensis Pic, 1945a: 6 (*Leptura*)
- montana montana*** Mulsant & Rey, 1863: 179 (*Leptura*) Çanakkale (Anatolia)
leuthneri Ganglbauer, 1886: 516 (*Leptura*)
nigerrima Pic, 1892i: 415 (*Leptura*)
pernigra Reitter, 1898c: 193 (*Leptura*)
semisanguinea Reitter, 1898c: 193 (*Leptura*)
bogasana Pic, 1932: 31 (*Leptura*)
claudiusi Pic, 1941b: 1 (*Leptura*)
juliae Pic, 1941b: 1 (*Leptura*)
mulsanti Pic, 1941b: 1 (*Leptura*)
victoris Pic, 1941b: 1 (*Leptura*)
- sanguinolenta*** Linnaeus, 1760: 196 (*Leptura*) İstanbul (Anatolia), Bursa
variabilis DeGeer, 1775: 137 (*Leptura*)
ignita Geoffroy, 1785: 89 (*Stenocorus*)
bonaeriensis Burmeister, 1865: 177 (*Leptura*)
sandoeensis Palm, 1953: 124 (*Leptura*)
- Genus ***Anoplodera*** Mulsant, 1839: 285
type species *Leptura sexguttata* Fabricius, 1775
- Subgenus ***Anoplodera*** Mulsant, 1839: 285
type species *Leptura sexguttata* Fabricius, 1775
- rufipes rufipes*** Schaller, 1783: 296 (*Leptura*) [HN] Kırklareli
atra Fabricius, 1775: 197 (*Leptura*) [NO]
villosa Schoenherr, 1817: 486 (*Leptura*)
fuscipes Mulsant, 1839: 287
bursensis Jureček, 1931: 124 (*Leptura*)
- rufipes ventralis*** Heyden, 1886: 85 [RN] Balıkesir, Bursa, Çanakkale (Anatolia)
rufiventris Tournier, 1872: 348 (*Leptura*) [HN]
medea Pic, 1909: 99 (*Leptura*)
- sexguttata*** Fabricius, 1775: 198 (*Leptura*) Kocaeli, İstanbul (Anatolia),
uddmanniana Harrer, 1784: 225 (*Leptura*) İstanbul (Thrace)
exclamationis Fabricius, 1793: 343 (*Leptura*)
semicolon Schrank, 1798: 694 (*Leptura*)
punctomaculata Marsham, 1802: 357 (*Leptura*)
biguttata Mulsant, 1839: 285
atrata Schilsky, 1889: 339
guttata Pic, 1898h: 4

pyrenaica Pic, 1917b: 5 (*Leptura*)
vastatorum Pic, 1917b: 5 (*Leptura*)

Genus ***Grammoptera*** Dejean, 1835: 356
type species *Leptura praeusta* Fabricius, 1787 (= *Leptura ustulata* Schaller, 1783)

Subgenus ***Grammoptera*** Dejean, 1835: 356
type species *Leptura praeusta* Fabricius, 1787 (= *Leptura ustulata* Schaller, 1783)
Fallaciomorpha Pic, 1900h: 37 type species *Grammoptera angustata* Pic, 1892j

abdominalis Stephens, 1831: 262 (*Leptura*) Kırklareli
variegata Germar, 1823: 522 (*Leptura*) [HN]
analıs Herrich-Schäffer, 1833: Heft 118 (6) (*Leptura*)
nigrescens Weise, 1884: 424

merkli Frivaldszky, 1884: 4 # *** Bursa

ruficornis ruficornis Fabricius, 1781: 247 (*Leptura*) [NP] İstanbul (Anatolia), Bursa,
İstanbul (Thrace)
rufipes Goeze, 1777: 501 (*Leptura*) [NO]
pumila Schaller, 1783: 299 (*Leptura*)
parisina Thunberg, 1784: 16 (*Leptura*)
laevis Herbst, 1784: 103 (*Leptura*)
clavipes Geoffroy, 1785: 87 (*Stenocorus*)
pallipes Stephens, 1831: 263 (*Leptura*)

ustulata ustulata Schaller, 1783: 298 (*Leptura*) Kırklareli
splendida Herbst, 1784: 103 (*Leptura*)
praeusta Fabricius, 1787: 159 (*Leptura*)
adusta Gmelin, 1790: 1872 (*Leptura*)
barrosi Pic, 1900c: 6
semirufescens Pic, 1947a: 4

Genus ***Judolia*** Mulsant, 1863: 496
type species *Leptura sexmaculata* Linnaeus, 1758

Subgenus ***Pachytodes*** Pic, 1891g: 65
type species *Leptura cerambyciformis* Schrank, 1781

cerambyciformis Schrank, 1781a: 154 (*Leptura*) Balıkesir, Bursa, Kırklareli
quadrinaculatus Scopoli, 1763: 47 (*Leptura*) [HN]
octomaculatus Schaller, 1783: 297 [wrongly as 299] (*Leptura*) [HN]
decempunctatus Olivier, 1800: 26 (*Leptura*)
sempunctatus Mulsant, 1839: 244 (*Pachyta*)
bisbistigma Pic, 1906c: 67 (*Leptura*)
salbachii Reitter, 1908: 216 (*Pachyta*)
anticeundulatus Pic, 1915a: 29 (*Leptura*)
bisquadristigma Pic, 1915a: 29 (*Leptura*)
digoniensis Pic, 1915a: 29 (*Leptura*)
urbisensis Pic, 1915a: 29 (*Leptura*)
valesiacus Pic, 1915a: 29 (*Leptura*)
multiinterruptus Pic, 1915a: 30 (*Leptura*)
beskidicus Pic, 1915c: 18 (*Leptura*)

humerifer Pic, 1915c: 18 (*Leptura*)
efasciatus Pic, 1916b: 9 (*Leptura*)
transylvanicus Pic, 1916b: 9 (*Leptura*)
parvonotatus Pic, 1916b: 10 (*Leptura*)
externeonotata Pic, 1923a: 13 (*Pachytodes*)
hoverlanus Roubal, 1937: 81
breveseperatus Pic, 1953: 9 (*Pachytodes*)
lateseparatus Pic, 1953: 9 (*Pachytodes*)

erraticus erythrurus Küster, 1848: 90 (*Pachyta*)
septemsignatus Küster, 1848a: 89 (*Pachyta*)
akbesianus Pic, 1898h: 6 (*Pachytodes*)
rosinae Pic, 1901: 11 (*Leptura*)
hungaricus Pic, 1913: 186 (*Leptura*)
kalavritanus Pic, 1913: 186 (*Leptura*)
rufopicalis Pic, 1913: 186 (*Leptura*)
rufonotatus Pic, 1913: 186 (*Leptura*)
testaceofasciatus Pic, 1913: 186 (*Leptura*)
anticonotatus Pic, 1914c: 13 (*Leptura*)
eibesanus Pic, 1914c: 13 (*Leptura*)
anticedivisus Pic, 1914c: 14 (*Leptura*)
unijunctus Pic, 1914c: 14 (*Leptura*)
subapicalis Pic, 1914c: 15 (*Leptura*)
quinquepunctatus Pic, 1915c: 18 (*Leptura*)
italicus Pic, 1916c: 4 (*Pachytodes*)
siculus Pic, 1916c: 4 (*Pachytodes*)
ragusai Pic, 1923b: 3 (*Pachytodes*)

İstanbul (Anatolia), Balıkesir,
Bilecik, Bursa, Çanakkale
(Anatolia), Yalova, Kırklareli,
İstanbul (Thrace)

Genus ***Leptura*** Linnaeus, 1758: 397
type species *Leptura quadrifasciata* Linnaeus, 1758

Subgenus ***Leptura*** Linnaeus, 1758: 397
type species *Leptura quadrifasciata* Linnaeus, 1758

Bothrioileptura Pesarini & Sabbadini, 2015: 27 type species *Leptura alticola* Gressitt, 1948
Rhytidoleptura Pesarini & Sabbadini, 2015: 29 type species *Leptura christinae* Pesarini, Rapuzzi
& Sabbadini, 2015 in Pesarini & Sabbadini, 2015

aurulenta Fabricius, 1793: 348
algerica Pic, 1898h: 6 (*Strangalia*)
viturati Pic, 1914b: 18 (*Strangalia*)
subreducta Pic, 1915a: 5 (*Strangalia*)
lugdunensis Pic, 1916a: 2: (*Strangalia*)
mesmini Pic, 1916c: 5 (*Strangalia*)
kitanensis Dayrem, 1922: 17 (*Strangalia*)

İstanbul (Anatolia), Balıkesir,
Kırklareli, İstanbul (Thrace)

quadrifasciata lederi Ganglbauer, 1882: 697
notatipennis Pic, 1897d: 5 (*Strangalia*)
suramensis Pic, 1915b: 5 (*Strangalia*)
caucasica Plavilstshikov, 1924: 226 (*Strangalia*)

İstanbul (Anatolia), Balıkesir

quadrifasciata quadrifasciata Linnaeus, 1758: 398
octomaculata DeGeer, 1775: 132
quadripustulata Fabricius, 1793: 345

İstanbul (Thrace)

apicalis Curtis, 1831: 362
apicata Stephens, 1839: 278
interrupta Heyden, 1877a: 397 (*Strangalia*)
guillemoti Desbrochers des Loges, 1895: 130 (*Stenura*)
melgunowi Jakobson, 1895: 523 (*Strangalia*)
mosquensis Pic, 1915b: 5 (*Strangalia*)
suramensis Pic, 1915b: 5 (*Strangalia*)
martialis Pic, 1941c: 1 (*Strangalia*)
bidivisa Schmidt, 1951: 13 (*Strangalia*)

Genus ***Pedostrangalia*** Sokolov, 1897: 461
type species *Pedostrangalia kassjanowi* Sokolov, 1897 (= *Leptura imberbis* Ménétriés, 1832)

Subgenus ***Neosphenalia*** Löbl, 2010: 60
type species *Leptura verticalis* Germar, 1822
Sphenalia K. Daniel, 1904b: 355 [HN] type species *Leptura verticalis* Germar, 1822

emmipoda Mulsant, 1863: 531 (*Strangalia*) # Balıkesir, Çanakkale (Anatolia)
insuturata Pic, 1891c: 15 (*Strangalia*)
subsignata Pic, 1901l: 235 (*Strangalia*)
adanensis Pic, 1917b: 6 (*Strangalia*)
tambei Pic, 1917b: 6 (*Strangalia*)
chehirensis Pic, 1933a: 6 (*Strangalia*)
perroudi Pic, 1933a: 6 (*Strangalia*)

verticalis Germar, 1822: 9 (*Leptura*) İstanbul (Thrace)
graeca Pic, 1901i: 52 (*Strangalia*)
taygetana K. Daniel, 1904b: 366 (*Leptura*)

Subgenus ***Pedostrangalia*** Sokolov, 1897: 461
type species *Pedostrangalia kassjanowi* Sokolov, 1897 (= *Leptura imberbis* Ménétriés, 1832)

revestita revestita Linnaeus, 1767: 638 (*Leptura*) Kirklareli
villica Fabricius, 1775: 196 (*Leptura*)
rubra Geoffroy, 1785: 89 (*Stenocorus*)
fuscicornis Marsham, 1802: 357 (*Leptura*)
ferruginea Mulsant, 1839: 254 (*Strangalia*)
fulvilabris Mulsant, 1839: 254 (*Strangalia*)
labiata Mulsant, 1839: 254 (*Strangalia*)
rufomarginata Mulsant, 1839: 254 (*Strangalia*)
vitticollis Mulsant, 1839: 254 (*Strangalia*)
discicollis W. G. H. Scriba, 1867: 32 (*Strangalia*)
diversipennis Pic, 1911d: 16 (*Leptura*)
bicoloraticeps Pic, 1911d: 17 (*Leptura*)
gabilloti Pic, 1911d: 18 (*Leptura*)
rufonotata Pic, 1914b: 18 (*Strangalia*)
lectorica Dayrem, 1916: 17 (*Leptura*)
marsolanensis Dayrem, 1916: 17 (*Leptura*)
brevenotaticollis Pic, 1933a: 14 (*Strangalia*)
barbanti Pic, 1933c: 5 (*Strangalia*)
schleicheri Pic, 1934b: 38 (*Leptura*)

Genus ***Pseudovadonia*** Lobanov, Danilevsky & Murzin, 1981: 787
type species *Leptura livida* Fabricius, 1777

livida livida Fabricius, 1777: 233 (*Leptura*)
pastinacae Panzer, 1795: 275 (*Leptura*)

Edirne, Kırklareli

livida setosa Danilevsky, 2013a: 29

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
İstanbul (Thrace)

Genus ***Rutpela*** Nakane & K. Ohbayashi, 1957: 48
type species *Leptura maculata* Poda von Neuhaus, 1761

maculata maculata Poda von Neuhaus, 1761: 37 (*Leptura*) Kocaeli, Sakarya, İstanbul
fasciata Scopoli, 1763: 54 (*Cerambyx*) (Anatolia), Balıkesir,
elongata DeGeer, 1775: 134 (*Leptura*) Bursa, Çanakkale
armata Herbst, 1784: 101 (*Leptura*) (Anatolia), Yalova,
scopoliana Laicharting, 1784: 145 (*Leptura*) Kırklareli, İstanbul
rubea Geoffroy, 1785: 86 (*Stenocorus*) (Thrace)
quinquemaculata Gmelin, 1790: 1868 (*Leptura*)
sinuata Fabricius, 1793: 347 (*Leptura*)
subspinosa Fabricius, 1793: 347 (*Leptura*)
calcarata Olivier, 1800: 14 (*Leptura*)
binotata Mulsant, 1839: 258 (*Strangalia*)
externepunctata Mulsant, 1839: 258 (*Strangalia*)
impunctata Mulsant, 1839: 258 (*Strangalia*)
punctatofasciata Mulsant, 1839: 258 (*Strangalia*)
undulata Mulsant, 1839: 259 (*Strangalia*)
dayremi Pic, 1903b: 4 (*Strangalia*)
alsatica Pic, 1906c: 67 (*Leptura*)
disconotata Pic, 1908b: 3 (*Leptura*)
bifenestrata Pic, 1933a: 2 (*Strangalia*)
nicodi Pic, 1933a: 6 (*Strangalia*)
wuenschi Roubal, 1937: 81 (*Strangalia*)
discoininterrupta Pic, 1945b: 13 (*Strangalia*)
dromensis Pic, 1945b: 13 (*Strangalia*)
pacifica Pic, 1945b: 13 (*Strangalia*)

manca Schaufuss, 1863: 121 (*Strangalia*)
nigrofasciata V. Petagna, 1792: 247 (*Leptura*) [HN]
nigricornis Stierlin, 1864: 153 (*Strangalia*)
kricheldorfii Wagner, 1928: 121 (*Leptura*)

İstanbul (Thrace)

Genus ***Stenurella*** Villiers, 1974: 214
type species *Leptura melanura* Linnaeus, 1758

Subgenus ***Nigrostenurella*** Özdikmen, 2013c: 525
type species *Leptura nigra* Linnaeus, 1758

nigra nigra Linnaeus, 1758: 398 (*Leptura*)
picea Geoffroy, 1785: 87 (*Stenocorus*)
varicollis Schaefer, 1932: 31 (*Stenura*)

Kocaeli, Kırklareli, İstanbul (Thrace)

giraudi Pic, 1946: 14 (*Strangalia*)

Subgenus **Priscostenurella** Özdikmen, 2013c: 516

type species *Leptura bifasciata* O.F.Müller, 1776

Eduardvivesia Zamoroka, Trócoli, Shparyk & Semaniuk, 2022: 131 type species *Leptura vaucheri* Bedel, 1900

Nigromacularia Zamoroka, Trócoli, Shparyk & Semaniuk, 2022 132 type species *Leptura septempunctata* Fabricius 1793

bifasciata intermedia Holzschuh, 2006: 219

Kırklareli, İstanbul (Thrace)

bifasciata limbiventris Reitter, 1898d: 21 (*Strangalia*)

Kocaeli, Balıkesir, Bilecik,
Bursa, Çanakkale
(Anatolia), Yalova

septempunctata latenigra Pic, 1915b: 5 (*Strangalia*) #

anatolica Heyrovský, 1961a: 45 (*Strangalia*)

roberti Pic, 1915a: 38 (*Leptura*)

afyoncayensis Şabanoğlu Şimşek, Özdemir Türkmen & Sert, 2024: 170 Kocaeli, İstanbul
(Anatolia), Balıkesir
Bilecik, Bursa,
Çanakkale
(Anatolia), Yalova,
Kırklareli, İstanbul
(Thrace)

Subgenus **Stenurella** Villiers, 1974: 214

type species *Leptura melanura* Linnaeus, 1758

melanura Linnaeus, 1758: 397 (*Leptura*)

suturanigra DeGeer, 1775: 138 (*Leptura*)

similis Herbst, 1784: 101 (*Leptura*)

diversiventris Dufour, 1843: 103 (*Leptura*)

georgiana Pic, 1891g: 63 (*Strangalia*)

latesaturata Pic, 1891g: 63 (*Strangalia*)

rubellata Reitter, 1901c: 77 (*Strangalia*)

melanurella Reitter, 1901c: 78 (*Strangalia*)

semicrassa Pic, 1901a: 58 (*Leptura*)

sennii Sama, 2002: 40

İstanbul (Anatolia), Balıkesir,
Bursa, Çanakkale (Anatolia),
Yalova, Kırklareli, İstanbul
(Thrace)

samai Rapuzzi, 1995: 617 #

Bursa, Kırklareli

Genus **Stictoleptura** Casey, 1924: 280

type species *Leptura cribripennis* LeConte, 1859

Subgenus **Aredolpona** Nakane & K. Ohbayashi, 1957: 50

type species *Leptura rubra* Linnaeus, 1758

Corymbia Gozis, 1886: 33 [HN] type species *Leptura rubra* Linnaeus, 1758

rubra rubra Linnaeus, 1758: 397 (*Leptura*)

testacea Linnaeus, 1760: 196 (*Leptura*)

umbellatarum Laicharting, 1784: 161 (*Leptura*)

rubrotestacea Illiger, 1805: 122 (*Leptura*)

occipitalis Mulsant, 1839: 268 (*Leptura*)

Kocaeli, Sakarya, İstanbul
(Anatolia), Kırklareli,
İstanbul (Thrace)

salzmanni Hubenthal, 1902: 291 (*Leptura*)
sattleri Bickhardt, 1902: 362 (*Leptura*)
maculiceps Gabriel, 1910: 556 (*Leptura*)
semimarginata Pic, 1916a: 22 (*Leptura*)
dufour Lecomte, 1926: 168 (*Leptura*)
planata Swaine & Hopping, 1928: 62 (*Anoplodera*)

Subgenus **Maculileptura** Danilevsky, 2014b: 274
type species: *Leptura maculicornis* DeGeer, 1775

maculicornis DeGeer, 1775: 139 (*Leptura*)
maculosa Gmelin, 1790: 1874 (*Leptura*)

Kırklareli

pallens Brullé, 1832: 264 (*Leptura*)

Kocaeli, İstanbul (Anatolia),
Çanakkale (Anatolia),
Kırklareli, İstanbul (Thrace)

Subgenus **Melanoleptura** Miroshnikov, 1998: 594
type species *Leptura scutellata* Fabricius, 1781

scutellata scutellata Fabricius, 1781: 247 (*Leptura*)
?carbonaria Scopoli, 1763: 56 (*Cerambyx*) [NO]
funerea Geoffroy, 1785: 89 (*Stenocorus*)
nigra V. Petagna, 1787: 17 (*Rhagium*)
chrysothyreos Schrank, 1798: 697 (*Leptura*)
ochraceipennis Pic, 1897e: 31 (*Leptura*)

İstanbul (Anatolia), Balıkesir,
Çanakkale (Anatolia),
Kırklareli, İstanbul (Thrace)

Subgenus **Paracorymbia** Miroshnikov, 1998: 587
type species *Leptura fulva* DeGeer, 1775

excisipes K. Daniel & J. Daniel, 1891: 6 (*Leptura*) #

Balıkesir

fulva DeGeer, 1775: 137 (*Leptura*)
lutescens Geoffroy, 1785: 87 (*Stenocorus*)
tomentosa Fabricius, 1793: 340 (*Leptura*)
affinis Marsham, 1802: 353 (*Leptura*)
apicalis Motschulsky, 1875: 142 (*Leptura*)
corsica Pic, 1894d: 206 (*Leptura*)
fulvoapicalis Plavilstshikov, 1932: 174 (*Leptura*)

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bursa, Çanakkale
(Anatolia), Yalova, Tekirdağ,
Kırklareli, İstanbul (Thrace)

pallidipennis Tournier, 1872: 346 (*Leptura*)
circassica K. Daniel & J. Daniel, 1891: 31 (*Leptura*)

İstanbul (Anatolia)

tonsa K. Daniel & J. Daniel, 1891: 31 (*Leptura*)
ustulata Ménétériés, 1832: 231 (*Leptura*) [HN]
menetriesi Ganglbauer, 1889d: 469 [RN] [NO]

Kocaeli, Yalova, İstanbul (Thrace)

Subgenus **Stictoleptura** Casey, 1924: 280
type species *Leptura cribripennis* LeConte, 1859

cordigera cordigera Fuessly, 1775: 14 (*Leptura*)

hastata Sulzer, 1776: 46 (*Leptura*)

lamed Geoffroy, 1785: 86 [HN] (*Stenocorus*)

immaculata Ragusa, 1889: 236 (*Leptura*)

auricollis Chobaut, 1895: cccx (*Leptura*)

inhastata Pic, 1902c: 8 (*Leptura*)

epipleuralis Schaefer, 1932: 31 (*Leptura*)

Kocaeli, Sakarya, İstanbul
(Anatolia), Balıkesir,
Bilecik, Bursa, Çanakkale
(Anatolia), Yalova, Edirne,
Tekirdağ, Kırklareli,
İstanbul (Thrace),
Çanakkale (Gelibolu)

rufa nigropicta Fairmaire, 1866: 278 (*Leptura*) # ***

Balıkesir

rufa rufa Brullé, 1832: 263 (*Leptura*)

İstanbul (Anatolia), Bilecik,
İstanbul (Thrace)

Genus ***Strangalia*** Dejean, 1835: 355

type species *Leptura luteicornis* Fabricius, 1775

Ophistomis J. Thomson, 1861: 154 type species *Ophistomis flavocinctus* J. Thomson, 1861

Ophistomis Gemminger, 1872: 2875 [unnecessary emendation]

Strangalina Aurivillius, 1912: 228 type species *Leptura attenuata* Linnaeus, 1758

Sulcatostrangalia K. Ohbayashi, 1961: 17 type species *Strangalia gracilis* Gressitt, 1935

attenuata Linnaeus, 1758: 398 (*Leptura*)

ucranica Laxman, 1770: 596

brunnescens Balbi, 1892: 49

obscuriventris Pic, 1901a: 59 (*Typocerus*)

imperfecta Gerhardt, 1910: 556 (*Leptura*)

grenieri Pic, 1912c: 3 (*Leptura*)

balcanica Pic, 1915b: 6 (*Typocerus*)

varcanica, Ohbayashi, 2023: 67 (misspelling)

İstanbul (Anatolia), Kırklareli,
İstanbul (Thrace)

Genus ***Vadonia*** Mulsant, 1863: 559

type species *Leptura unipunctata* Fabricius, 1787

Subgenus ***Sinemaculipenna*** Özdikmen, 2025c: 2989

type species *Vadonia instigmata* Pic, 1889

bitlisiensis Chevrolat, 1882: 59 #

Bilecik

cribricollis Pic, 1889a: 20 [mispaginated: 5] (*Leptura*)

bistigmata Pic, 1890b: clxxvi

armeniaca Pic, 1903b: 4 (*Leptura*)

Subgenus ***Vadonia*** Mulsant, 1863: 559

type species *Leptura unipunctata* Fabricius, 1787

Neovadonia Kaszab, 1938: 151 type species *Leptura unipunctata* Fabricius, 1787

bolognai Sama, 1982: 207 # ***

Kocaeli

grandicollis grandicollis Mulsant & Rey, 1863: 182 #

İstanbul (Anatolia),
İstanbul (Thrace)

bisignata Brullé, 1832: 264 (*Leptura*) [HN]

inapicalis Pic, 1897e: 31 (*Leptura*) [DA]

- moesiaca** K. Daniel & J. Daniel, 1891: 6 (*Leptura*)
bipunctata Kraatz, 1859: 97 [HN]
beali Pic, 1926a: 13 [DA] İstanbul (Anatolia), Balıkesir,
Edirne, Kırklareli
- unipunctata unipunctata** Fabricius, 1787: 157 (*Leptura*) [NP] İstanbul (Anatolia),
pilosa Forster, 1771: 44 (*Leptura*) [NO] Bilecik, Edirne,
uninstigmata Pic, 1891c: 9 (*Leptura*) İstanbul (Thrace)
obscorepilosa Pic, 1892l: lxxxiv (*Leptura*)

Subfamily NECYDALINAE Latreille, 1825

Tribe Necydalini Latreille, 1825

Genus **Necydalis** Linnaeus, 1758: 421
type species *Necydalis major* Linnaeus, 1758

Subgenus **Necydalis** Linnaeus, 1758: 421
type species *Necydalis major* Linnaeus, 1758
Gymnopterion Schrank, 1798: 373 type species *Necydalis major* Linnaeus, 1758

- major** Linnaeus, 1758: 421 Kırklareli
ichneumonea DeGeer, 1775: 148
abbreviata Fabricius, 1775: 199 (*Leptura*)
majus Schrank, 1798: 688 (*Gymnopterion*)
populi Büttner, 1818: 245 (*Molorchus*)
salicis Mulsant, 1839: 112
duponti Mulsant, 1839: 299 (*Molorchus*)
xantha Semenov, 1899a: 132
subnotata Pic, 1941b: 1
altaica Pic, 1941b: 2
rufiventris Pic, 1941b: 2
aino Kusama, 1974: 54
- ulmi** Chevrolat, 1838: 76 (*Molorchus*) [NP] Kırklareli
annulata L. Petagna, 1819: 19 (*Molorchus*) [NO]
panzeri Harold, 1876: 174
mesembrina Plavilstshikov, 1936: 467
hadullai Szallies, 1994: 260

Subfamily SAPHANINAE Gistel, 1848

Tribe Saphanini Gistel, 1848

Genus **Saphanus** Audinet-Serville, 1834a: 81
type species *Callidium spinosum* Fabricius, 1793 (= *Callidium piceum* Laicharting, 1784)

- kadleci** Rapuzzi & Sama, 2014b: 4 # *** Kocaeli, Sakarya
- piceus ganglbaueri** Brancsik, 1886: 71 Kırklareli

Tribe **Anisarthrini** Mamaev & Danilevsky, 1973

Genus **Alocerus** Mulsant, 1862: 127

type species *Alocerus fulvus* Mulsant, 1862 (= *Callidium moesiacum* Frivaldszky von Frivald, 1837)

Cyamophthalmus Kraatz, 1863: 99 type species *Cyamophthalmus ferrugineus* Kraatz, 1863 = *Callidium moesiacus* Frivaldszky von Frivald, 1837)

moesiacus Frivaldszky von Frivald, 1837: 177 (*Callidium*)

Balıkesir, Kırklareli

fulvus Mulsant, 1862: 128

ferrugineus Kraatz, 1863: 100 (*Cyamophthalmus*)

nitidus Fairmaire, 1864b: 340 (*Cyamophthalmus*)

syriacus Chevrolat, 1882: 59 (*Smodicum*)

Subfamily **ASEMINAE** J. Thomson, 1861

Tribe **Asemini** J. Thomson, 1861

Genus **Arhopalus** Audinet-Serville, 1834a: 77

type species *Cerambyx rusticus* Linnaeus, 1758

Criocephalum Dejean, 1835: 328 type species *Cerambyx rusticus* Linnaeus, 1758

Criocephalus Mulsant, 1839: 63 type species *Cerambyx rusticus* Linnaeus, 1758

Hylescopus Gistel, 1856: 376 [unnecessary substitute name]

ferus Mulsant, 1839: 64 (*Criocephalus*)

Balıkesir, Bilecik, Kırklareli

polonicus Motschulsky, 1845: 88 (*Criocephalum*)

pinetorum Wollaston, 1863: 103 N: CI

epibatus Schiödt, 1864: 521 (*Criocephalum*)

rusticus Linnaeus, 1758: 395 (*Cerambyx*)

İstanbul (Anatolia), Balıkesir,

tristis Fabricius, 1787: 154 (*Callidium*)

Çanakkale (Anatolia), İstanbul

pachymerus Mulsant, 1839: 64 (*Criocephalus*)

(Thrace), Çanakkale (Gelibolu)

lugubris Gmelin, 1790: 1847 (*Cerambyx*)

coriaceus Motschulsky, 1845: 89 (*Criocephalum*)

hispanicus Sharp, 1905a: 157 (*Criocephalus*)

Genus **Cephalocrius** Sharp, 1905a: 148

type species *Criocephalus syriacus* Reitter, 1895

syriacus Reitter, 1895a: 86 (*Criocephalus*) #

Balıkesir

Subfamily **SPONDYLIDINAE** Audinet-Serville, 1832

Tribe **Spondylidini** Audinet-Serville, 1832

Genus **Spondylis** Fabricius, 1775: 159

type species *Attelabus buprestoides* Linnaeus, 1758

Sphondyla Illiger, 1804: 115 [unjustified emendation]

Sphondylis Gistel, 1848b: 129 [unnecessary substitute name]

buprestoides buprestoides Linnaeus, 1758: 388 (*Attelabus*)
maxillosus DeGeer, 1775: 84 (*Cerambyx*)
elongatum Latreille, 1829: 107
sinensis Nonfried, 1892b: 92
zwergii Bodemeyer, 1927: 43

İstanbul (Anatolia),
Balıkesir, Bursa,
İstanbul (Thrace)

Subfamily CERAMBYCINAE Latreille, 1802

Tribe Hesperophanini Mulsant, 1839

Subtribe Hesperophanina Mulsant, 1839

Genus ***Hesperophanes*** Dejean, 1835: 328
type species *Callidium sericeum* Fabricius, 1787

sericeus Fabricius, 1787: 152 (*Callidium*)
latreillei Brullé, 1832: 256 (*Callidium*)
rotundicollis P. H. Lucas, 1842: 185

İstanbul (Anatolia), Balıkesir,
İstanbul (Thrace)

Genus ***Stromatium*** Audinet-Serville, 1834a: 80
type species *Callidium barbatum* Fabricius, 1775
Solenophorus Mulsant, 1839: 65 type species *Callidium strepens* Fabricius, 1798 (= *Callidium unicolor* Olivier, 1800)

auratum Böber, 1793: 135 (*Saperda*)
fulvum Villers, 1789: 256 (*Cerambyx*) [HN]
strepens Fabricius, 1798: 150 (*Callidium*)
unicolor Olivier, 1800: no. 70: 58 (*Callidium*)
pallidum Zubkov, 1833: 336 (*Callidium*)
inermis Tournier, 1872: 260
platifemur Chevrolat, 1882: 57 (*Hesperophanes*)

Sakarya, İstanbul (Anatolia),
Balıkesir, Bursa, Çanakkale
(Anatolia), Edirne, Kırklareli,
İstanbul (Thrace)

Genus ***Trichoferus*** Wollaston, 1854: 427
type species *Trichoferus senex* Wollaston, 1854 (= *Trichoferus fasciculatus senex* Wollaston, 1854)

Subgenus ***Trichoferus*** Wollaston, 1854: 427
type species *Trichoferus senex* Wollaston, 1854 (= *Trichoferus fasciculatus senex* Wollaston, 1854)
Hesperandrius Reitter, 1913a: 45 type species *Callidium griseum* Fabricius, 1793

fasciculatus fasciculatus Faldermann, 1837: 266 (*Hesperophanes*)
pulverulentus Erichson, 1841: 188 (*Callidium*)
affinis P. H. Lucas, 1842: 186 (*Hesperophanes*)
elongatulus Pic, 1896a: 35 (*Hesperophanes*)
pubescens Pic, 1898a: 9 (*Hesperophanes*)
subnudus Dayrem, 1928b: 67 (*Hesperophanes*)

Balıkesir, Bursa,
Çanakkale (Anatolia)

griseus Fabricius, 1793: 325 (*Callidium*)
bimaculatus Bilberg, 1817: 194 (*Clytus*)

İstanbul (Thrace)

tomentosus P. H. Lucas, 1842: 186 (*Hesperophanes*)
decolor Pic, 1937: 14 (*Hesperophanes*)

holosericeus Rossi, 1790: 153 (*Callidium*) İstanbul (Anatolia), İstanbul (Thrace)
cinereus Villers, 1789: 256 (*Cerambyx*) [HN]
nebulosus Olivier, 1790: 257 (*Callidium*)
gayi Winkler, 1929: 1143 [RN] [not RN by Plavilstshiov, 1921a]

Tribe **Phoracanthini** Newman, 1840

Genus **Cordylomera** Audinet-Serville, 1834a: 23
type species *Cordylomera nitipennis* Audinet-Serville, 1834 = *Cerambyx spinicornis* Fabricius, 1775
Acanthinomomus Hope, 1835: 107 type species *Cerambyx spinicornis* Fabricius, 1775

spinicornis Fabricius, 1775: 167 (*Cerambyx*) Kocaeli
torrida Olivier, 1795: 31 (*Cerambyx*)
testacea Buquet, 1844: 233
suturalis Chevrolat, 1858: 242
sansibarica Kolbe, 1893: 248
angolensis Veiga-Ferreira, 1971: 7

Tribe **Cerambycini** Latreille, 1802

Genus **Cerambyx** Linnaeus, 1758: 388
type species *Cerambyx cerdo* Linnaeus, 1758

Subgenus **Cerambyx** Linnaeus, 1758: 388
type species *Cerambyx cerdo* Linnaeus, 1758
Hamaticherus Dejean, 1821: 105 type species *Cerambyx heros* Scopoli, 1763 (= *Cerambyx cerdo* Linnaeus, 1758)
Hammatocerus Gistel, 1848b: 130 [unjustified emendation] [HN]

cerdo acuminatus Motschulsky, 1853: 79 Kocaeli, Sakarya, İstanbul (Anatolia), Balıkesir, Bilecik, Bursa, Çanakkale (Anatolia)

cerdo cerdo Linnaeus, 1758: 392 Kırklareli, İstanbul (Thrace)
heros Scopoli, 1763: 31
laevicollis Heyrovsky, 1955: 156 (as *morpha* - unavailable name)
laevicollis Vartanis, 2024e: 1144 (“the area of Třeboňsk, in the Czech Republic”) - unavailable name.

dux Faldermann, 1837: 264 (*Hammaticherus*) Bilecik, Bursa, Kırklareli, İstanbul (Thrace)
orientalis Küster, 1845: 45 (*Hammaticherus*)
thirkii Küster, 1845: 47 (*Hammaticherus*)
intricatus Fairmaire, 1848: 167 (*Hammaticherus*)
nodosus Mulsant & Rey, 1863: 144

miles Bonelli, 1812: 178 Edirne, İstanbul (Thrace)
militaris Latreille, 1829: 116 (*Hamaticherus*)

nodulosus nodulosus Germar, 1817: 220 İstanbul (Anatolia), Bursa,
nodicornis Küster, 1845: 43 (*Hammaticherus*) İstanbul (Thrace)

welensii welensii Küster, 1845: 44 (*Hammaticherus*) Sakarya, İstanbul (Anatolia),
velutinus Brullé, 1832: 252 [HN] Balıkesir, Bursa, İstanbul
(Thrace)

Subgenus ***Microcerambyx*** Mikšić & Georgijevic, 1973: 22
type species *Cerambyx scopolii* Fuessly, 1775

scopolii scopolii Fuessly, 1775: 12 Sakarya, İstanbul (Anatolia),
piceus Geoffroy, 1785: 74 Tekirdağ, Kırklareli, İstanbul
helveticus Stierlin, 1878a: 442 (Thrace)

Tribe **Compsoicerini** J. Thomson, 1864

Genus ***Rosalia*** Audinet-Serville, 1834b: 561
type species *Cerambyx alpinus* Linnaeus, 1758

Subgenus ***Rosalia*** Audinet-Serville, 1834b: 561
type species *Cerambyx alpinus* Linnaeus, 1758
Charmides Gistel, 1848a: [2] [unnecessary substitute name]

alpina Linnaeus, 1758: 392 (*Cerambyx*) Sakarya, İstanbul (Anatolia),
pilosa Poda von Neuhaus, 1761: 32 (*Cerambyx*) Balıkesir, Çanakkale (Anatolia),
croissandeau Pic, 1891c: 22 Kırklareli, İstanbul (Thrace)
multimaculata T. Pic, 1900a: 12
obliterata T. Pic, 1900a: 12
bifasciata Reitter, 1900a: 131
connexa Reitter, 1900a: 131
geminata Reitter, 1900a: 131
interrupta Reitter, 1900a: 131
prolongata Reitter, 1900a: 131
parvonotata Reitter, 1900a: 132
quadripunctata Reitter, 1901b: 202
quadrinaculata Ragusa, 1905: 8
gelineki Zoufal, 1906: 264
kyselyi Zoufal, 1906: 264
transsylvanica Pic, 1908c: 20
latemaculata T. Pic, 1908: 33
v-notata T. Pic, 1908: 33
brancsiki Pic, 1911b: 5
cepelaki Pic, 1933a: 6
triformis Roubal, 1937: 81
bystricensis Roubal, 1937: 82
korbeli Roubal, 1937: 82
uzunkumensis Podaný, 1963: 10

Tribe **Purpuricenini** J. Thomson, 1861

Genus **Calchaenesthes** Kraatz, 1863: 97

type species *Callidium oblongomaculatum* Guérin-Ménéville, 1844

Calchaenistes Schaum, 1862 [nomen oblitum] type species *Callidium nogeli* Frivaldszky von Frivald, 1845

Hysginois Truqui in Kraatz, 1863: 98 type species *Hysginois quadrigeminus* Truqui in Kraatz, 1863: 98 (= *Callidium oblongomaculatum* Guérin-Ménéville, 1844)

oblongomaculata Guérin-Ménéville, 1844: 234 (*Callidium*) İstanbul (Thrace)
nogeli Frivaldszky von Frivald, 1845: 183 (*Callidium*)
quadrimaculata Pic, 1912a: 13
subjuncta Pic, 1945a: 6

Genus **Purpuricen** Dejean, 1821: 105

type species *Cerambyx kaehleri* Linnaeus, 1758

Subgenus **Basiatropansexus** Özdikmen, 2025g: 2789

type species *Purpuricen barbarus* Lucas, 1842

dalmatinus Sturm, 1843: 353 Bilecik
biinterruptus Pic, 1901c: 11
disjunctus Pic, 1916a: 13
bonvouloiri Pic, 1945a: 6
hirsutus Heyrovský, 1969: 34

inhumeralis Pic, 1891c: 24 # Balıkesir, Çanakkale (Anatolia), Edirne

interscapillatus interscapillatus Plavilstshikov, 1937a: 247 [RN] # Bursa
humeralis Pic, 1891c: 23 ["Asie Mineure"] [HN]

Subgenus **Purpuricen** Dejean, 1821: 105

type species *Cerambyx kaehleri* Linnaeus, 1758

Acanthoptera Latreille, 1829: 114 type species *Cerambyx budensis* Götze, 1783

Sternoplistes Guérin-Ménéville, 1844: 224 type species *Purpuricen temminckii* Guérin-Ménéville, 1844

Cyclodera White, 1846: 510 type species *Cyclodera quadrinotata* White, 1846

Hamadrias Gistel, 1848b: 130 [unnecessary substitute name]

Philagathes J. Thomson, 1864: 196 type species *Philagathes laetus* J. Thomson, 1864

Porphyrocenus Reitter, 1913a: 34 type species *Purpuricen spectabilis* Motschulsky, 1858

budensis Götze, 1783: 72 (*Cerambyx*) Kocaeli, İstanbul
ungaricus Herbst, 1784: 90 (*Lamia*) (Anatolia), Balıkesir,
wredii Fischer von Waldheim, 1823: tab. 49, fig. 2 [1824: 238] Bursa, Çanakkale
affinis Brullé, 1832: 251 (Anatolia), Edirne,
İstanbul (Thrace)

kaehleri rossicus Danilevsky, 2019: 86 İstanbul (Anatolia), Kırklareli,
İstanbul (Thrace)

neocaucasicus Rapuzzi & Sama, 2014a: 156 # *** Bursa

renyvonae latefasciatus Özdikmen & Tezcan, 2020a: 537 # *** Kocaeli, İstanbul
(Anatolia), Balıkesir,
Bursa

renyvonae renyvona Sláma, 2001: 225 Edirne, Kırklareli, İstanbul (Thrace)

skypetarum zinaidae Lazarev, 2020a # *** Kocaeli, İstanbul (Anatolia)

Tribe **Callichromatini** Swainson & Shuckard, 1840

Genus **Aromia** Audinet-Serville, 1834a: 559
type species *Cerambyx moschatus* Linnaeus, 1758

Subgenus **Aromia** Audinet-Serville, 1834a: 559
type species *Cerambyx moschatus* Linnaeus, 1758
Terambus Gistel, 1848a: [2] [unnecessary substitute name]

moschata moschata Linnaeus, 1758: 391 (*Cerambyx*) Kocaeli, Sakarya,
İstanbul (Anatolia),
odorata DeGeer, 1775: 63 (*Cerambyx*) Balıkesir,
chlorophana Fischer von Waldheim, 1823: pl. xlviii [= 1824: 237] (*Cerambyx*) Bursa, İstanbul
alata Costa, 1855: 25 (Thrace)
auctumnalis Westhoff, 1882: 241
thea Reitter, 1894b: 306
cupricollis Pic, 1941a: 5
perroudi Pic, 1941a: 5

thoracica Fischer von Waldheim, 1823: tab. 48, figs. 3, 4 [1824: 236] (*Cerambyx*)
melancholica Reitter, 1895b: 210 Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
İstanbul (Thrace)

Tribe **Graciliini** Mulsant, 1839

Genus **Gracilia** Audinet-Serville, 1834a: 81
type species *Callidium pygmaeum* Fabricius, 1793 (= *Saperda minuta* Fabricius, 1781)
Nothrus Haldeman, 1847: 43 type species *Nothrus fuscus* Haldeman, 1847 (= *Saperda minuta* Fabricius, 1781)
Aphelocera Gistel, 1848a: [2] type species *Callidium vini* Panzer, 1799 (= *Saperda minuta* Fabricius, 1781)
Oesyophila Bedel, 1894: 156 [RN] type species *Callidium pygmaeum* Fabricius, 1793 (= *Saperda minuta* Fabricius, 1781)

minuta Fabricius, 1781: 235 (*Saperda*) İstanbul (Anatolia), Balıkesir,
picea Fabricius, 1793: 317 (*Saperda*) Çanakkale (Anatolia), İstanbul
pygmaea Fabricius, 1793: 323 (*Callidium*) (Thrace)
vini Panzer, 1799: 10 (*Callidium*)
fusca Haldeman, 1847: 43 (*Nothrus*)
approximata Fairmaire, 1883: clx
obliquata Horn, 1885: 174
albanica Csiki, 1931: 278

Genus **Penichroa** Stephens, 1839: 270

type species *Callidium fasciatum* Stephens, 1839

Liagrica Costa, 1855: 59 type species *Callidium timidum* Ménétériés, 1832 (= *Callidium fasciatum* Stephens, 1831)

Exilia Mulsant, 1862: 195 type species *Callidium timidum* Ménétériés, 1832 (= *Callidium fasciatum* Stephens, 1831)

Graecoeme Adlbauer, 2006: 379 type species *Graecoeme eggeri* Adlbauer, 2006

fasciata Stephens, 1831: 250 (*Callidium*)

Bursa, İstanbul (Thrace)

timida Ménétériés, 1832: 228 (*Callidium*)

bipunctata Zubkov, 1833: 336 (*Callidium*)

fasciolata Krynicki, 1834: 170 (*Callidium*)

brunnea Tournier, 1872: 280 (*Exilia*)

lugubris Ragusa, 1884: 333 (*Exilia*)

champlaini Knull, 1941: 695 (*Tylonotus*)

fagnezi Pic, 1945a: 6

eggeri Adlbauer, 2006: 381 (*Graecoeme*)

Tribe **Obrini** Mulsant, 1839

Genus **Obrium** Dejean, 1821: 110

type species *Cerambyx cantharinus* Linnaeus, 1767

Phyton Newman, 1840a: 19 type species *Phyton limum* Newman, 1840 (= *Cerambyx maculatus* Olivier, 1800)

Diozodes Haldeman, 1847: 42 type species *Callidium pallidum* Say, 1824 (= *Cerambyx maculatus* Olivier, 1800)

brunneum Fabricius, 1793: 316 (*Saperda*)

Bilecik, Bursa

caucasicum Tournier, 1872: 279

cantharinum cantharinum Linnaeus, 1767: 637 (*Cerambyx*)

İstanbul (Thrace)

ferrugineum Fabricius, 1781: 235 (*Saperda*)

fuscicorne Gmelin, 1790: 1837 (*Cerambyx*)

Tribe **Certallini** Fairmaire, 1864

Genus **Certallum** Dejean, 1821: 111

type species *Saperda ruficollis* Fabricius, 1781 (= *Cerambyx ebulinus* Linnaeus, 1767)

Cartallum Audinet-Serville, 1834a: 94 [unjustified emendation]

ebulinum Linnaeus, 1767: 637 (*Cerambyx*)

İstanbul (Anatolia), Balıkesir,

ruficolle Fabricius, 1781: 236 (*Callidium*)

Bilecik, Bursa, Çanakkale

monspeliense Gmelin, 1790: 1842 (*Cerambyx*)

(Anatolia), İstanbul (Thrace)

tricolor Chevrolat, 1882: 57 (*Obrium*)

nigricolle Pic, 1891c: 17 (*Cartallum*)

Tribe **Hylotruperini** Rose, 1983

Genus **Hylotrupes** Audinet-Serville, 1834a: 77

type species *Cerambyx bajulus* Linnaeus, 1758

Hylotrupes Gistel, 1856: 376 [unnecessary substitute name]

bajulus Linnaeus, 1758: 396 (*Cerambyx*)
caudatus DeGeer, 1775: 86 (*Cerambyx*)
linneianus Laicharting, 1784: 69 (*Callidium*)
quadripunctatus Geoffroy, 1785: 82 (*Leptura*)
similis Marsham, 1802: 335 (*Cerambyx*)
affinis Savenius, 1825: 63 (*Callidium*)
puellus A. Villa & G. B. Villa, 1833: 36 (*Callidium*)
lividus Mulsant, 1839: 56
bullatus Haldeman, 1847: 36
koziorowiczi Desbrochers des Loges, 1873: 429
syriacus Théry, 1896: 110
inaequalis Casey, 1924: 233
latus Casey, 1924: 233
incertus Casey, 1924: 234
nematocerus Casey, 1924: 234
puncticollis Casey, 1924: 234
minutus Casey, 1924: 235
pedicellatus Casey, 1924: 235
theresae Pic, 1924: 26

Sakarya, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
İstanbul (Thrace)

Tribe **Callidiini** Kirby, 1837

Genus ***Callidium*** Fabricius, 1775: 187
type species *Cerambyx violaceus* Linnaeus, 1758

Subgenus ***Callidostola*** Reitter, 1913a: 37
type species *Cerambyx aeneum* DeGeer, 1775
Calliopedia Binder, 1915: 186 type species *Rhopalopus reitteri* Binder, 1915 (= *Cerambyx aeneus* DeGeer, 1775)

aeneum aeneum DeGeer, 1775: 89 (*Cerambyx*)
variabile Fabricius, 1775: 188
cognatum Laicharting, 1784: 58
aurichalceum Gmelin, 1790: 1857 (*Cerambyx*)
dilatatum Paykull, 1800: 91
viride Schoenherr, 1817a: 459
venosum Eschscholtz, 1818: 483
viridans Gmelin, 1790: 1846 (*Cerambyx*)
semitestaceum Pic, 1945b: 6

Balıkesir

Genus ***Lioderina*** Ganglbauer, 1886: 517
type species *Callidium lineare* Hampe, 1871

linearis Hampe, 1871: 335 (*Callidium*)
henschi Puton, 1888: 23 (*Callidium*)

Bursa

Genus ***Phymatodes*** Mulsant, 1839: 47
type species *Cerambyx variabilis* Linnaeus, 1760 (= *Cerambyx testaceus* Linnaeus, 1758)

Subgenus ***Melasmetus*** Reitter, 1913a: 39
type species *Callidium femorale* Ménétériés, 1832

femoralis demelti Heyrovský, 1962a: 41 # ***

İstanbul (Anatolia), Bursa,
İstanbul (Thrace)

Subgenus *Phymatodellus* Reitter, 1913a: 40

type species *Callidium rufipes* Fabricius, 1777

Phymatodina Casey, 1912: 281 type species *Phymatodes nitidus* LeConte, 1874

Microcallidium Casey, 1912: 283 type species *Callidium amoenus* Say, 1824: 413

Pseudopocilium Planet, 1924: 226 type species *Callidium rufipes* Fabricius, 1777

rufipes rufipes Fabricius, 1777: 232 (*Callidium*)

Sakarya

coeruleus Goeze, 1777: 507 (*Leptura*)

cyaneus Geoffroy, 1785: 82 (*Leptura*)

cyanochrysos Gmelin, 1790: 1846 (*Cerambyx*)

erythropus Gmelin, 1790: 1846 (*Cerambyx*)

gallicus Gmelin, 1790: 1877 (*Leptura*)

amethystinus Fabricius, 1793: 321 (*Callidium*)

Subgenus *Phymatoderus* Reitter, 1913a: 39

type species *Callidium pusillum* Fabricius, 1787

Reitteroderus Sama, 1991: 124 [unnecessary substitute name]

lividus Rossi, 1794: 98 (*Callidium*)

Sakarya, İstanbul (Anatolia), Balıkesir

melancholicus Fabricius, 1798: 151 (*Callidium*)

brevicollis Dalman, 1817b: 191 (*Callidium*)

thoracicus Comolli, 1837: 44 (*Callidium*)

asperipennis Fairmaire, 1879: 217 (*Callidium*)

tristis Pic, 1891c: 20

pusillus pusillus Fabricius, 1787: 155 (*Callidium*)

İstanbul (Anatolia),
İstanbul (Thrace)

luridus Olivier, 1800: 23 (*Callidium*)

abdominalis Bonelli, 1812: 180 (*Callidium*)

humeralis Comolli, 1837: 44 (*Callidium*)

Subgenus *Phymatodes* Mulsant, 1839: 47

type species *Cerambyx variabilis* Linnaeus, 1760 (= *Cerambyx testaceus* Linnaeus, 1758)

Merium Kirby, 1837: 172 type species *Merium testaceus* Linnaeus, 1758 [suppressed]

testaceus Linnaeus, 1758: 396 (*Cerambyx*)

İstanbul (Anatolia), Balıkesir,
Bursa, Çanakkale (Anatolia),
Edirne, Kırklareli, İstanbul
(Thrace)

fennicus Linnaeus, 1758: 396 (*Cerambyx*)

variabilis Linnaeus, 1760: 192 (*Cerambyx*)

crassipes Goeze, 1777: 507 (*Leptura*)

praeustus Fabricius, 1782: 500 (*Callidium*)

femoratus Geoffroy, 1785: 83 (*Leptura*)

fallax Villers, 1789: 276 (*Leptura*)

violaceus Rossi, 1790: 154 (*Callidium*)

deustus Gmelin, 1790: 1850 (*Cerambyx*)

italicus Gmelin, 1790: 1851 (*Cerambyx*)

carniolicus Gmelin, 1790: 1858 (*Cerambyx*)

luridus Paykull, 1800: 87 (*Callidium*) [HN]

melanocephalus Ponza, 1805: 71 (*Cerambyx*)

nigrinus Mulsant, 1839: 47

nigricollis Mulsant, 1839: 49

similaris Küster, 1844: 54 (*Callidium*)
ventralis Haldeman, 1847: 375 (*Tessaropa*)
analís L. Redtenbacher, 1849: 481
rufipes Costa, 1855: 34
sellae Kraatz, 1868: 338 (*Callidium*)
dimidiatipennis Chevrolat, 1882: 58
fulvipennis Reitter, 1901a: 101 (*Phymatodes*)
barbarorus Pic, 1917a: 5
elongatus Iablokoff-Khnzorian, 1953: 60

Subgenus **Poecilium** Fairmaire, 1864a: 134
type species *Leptura alni* Linnaeus, 1767

alni alni Linnaeus, 1767: 639 (*Leptura*)
biarcuatus Piller & Mitterpacher, 1783: 112 (*Leptura*)
turcicus Geoffroy, 1785: 83 (*Leptura*)
infuscatus Chevrolat, 1866: 107 (*Poecilium*)
nitidus Chevrolat, 1882: 58 (*Callidium*)
apicalis Pic, 1915a: 1 (*Poecilium*)
fauconneti Pic, 1915a: 1 (*Poecilium*)
notaticollis Pic, 1945a: 6 (*Poecilium*)

Sakarya, İstanbul (Anatolia),
İstanbul (Thrace)

Genus **Pyrrhidium** Fairmaire, 1864a: 133
type species *Cerambyx sanguineus* Linnaeus, 1758

sanguineum Linnaeus, 1758: 396 (*Cerambyx*)

İstanbul (Thrace)

Genus **Ropalopus** Mulsant, 1839: 40
type species *Callidium clavipes* Fabricius, 1775

Subgenus **Ropalopus** Mulsant, 1839: 40
type species *Callidium clavipes* Fabricius, 1775
Rhopalopus Agassiz, 1846: 325 [unjustified emendation]
Euryoptera Horn, 1860: 571 type species *Euryoptera sanguinicollis* Horn, 1860

clavipes Fabricius, 1775: 188 (*Callidium*)
nigricans Gmelin, 1790: 1857 (*Cerambyx*)
nigroplanus DeGeer, 1775: 88 (*Cerambyx*)
viduus Geoffroy, 1785: 82 (*Leptura*)

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Yalova, Edirne, Kırklareli,
İstanbul (Thrace)

femoratus Linnaeus, 1758: 395 (*Cerambyx*)
punctuosus Geoffroy, 1785: 83 (*Leptura*)
castaneipennis Roubal, 1934a: 43

Edirne

macropus Germar, 1824: 514 (*Callidium*)
pilicollis C. G. Thomson, 1866: 29 (*Callidium*)
caucasicus Desbrochers des Loges, 1873: cxxxvi (*Callidium*)

Kocaeli, İstanbul (Anatolia),
Kırklareli, İstanbul (Thrace)

ungaricus insubricus Germar, 1823: 154 (*Callidium*)
viridipennis Pic, 1926d: 6 (*Rhopalopus*)

İstanbul (Anatolia),
İstanbul (Thrace)

Genus **Semanotus** Mulsant, 1839: 54

type species *Cerambyx undatus* Linnaeus, 1758

Sympiezocera P. H. Lucas, 1852: cvi type species *Sympiezocera laurasii* P. H. Lucas, 1852

Xenodorum Marseul, 1856: 48 type species *Xenodorum bonvouloiri* Marseul, 1856 (= *Sympiezocera laurasii* LeConte, 1851)

Anocomis Casey, 1912: 271 type species *Callidium ligneum* Fabricius, 1787

Hemicallidium Casey, 1912: 273 type species *Physocnenum amethystinum* LeConte, 1853

russicus russicus Fabricius, 1777: 232 (*Callidium*)

Balıkesir

Tribe **Anaglyptini** Lacordaire, 1868

Genus **Anaglyptus** Mulsant, 1839: 91

type species *Leptura mystica* Linnaeus, 1758

Subgenus **Anaglyptus** Mulsant, 1839: 91

type species *Leptura mystica* Linnaeus, 1758

arabicus Küster, 1847a: 95 (*Clytus*) #

İstanbul (Anatolia)

deyrollei Tournier, 1872: 277

mysticus Linnaeus, 1758: 398 (*Leptura*)

Kırklareli

quadricolor Scopoli, 1763: 56 (*Cerambyx*)

albofasciatus DeGeer, 1775: 82 (*Cerambyx*)

hieroglyphicus Herbst, 1784: 99 (*Callidium*)

litteratus Gmelin, 1790: 1857 (*Cerambyx*)

ruficornis Pic, 1908a: 35

bequarti Pic, 1910b: 3

obscurissimus Pic, 1901a: 59 # ***

Kocaeli

subimpressus Pic, 1901b: 9 [“Trébizonde”]

amasinus Pic, 1910a: 10 [“Amasya”]

anatolicus Demelt, 1970: 32 [“Samsun: Kavak”]

Genus **Paraclytus** Bates, 1884: 234

type species *Paraclytus excultus* Bates, 1884

sexguttatus Adams, 1817: 308 (*Clytus*)

Bursa, Kırklareli

bruckii Kraatz, 1864: 389 (*Clytus*)

caucasicus Motschulsky, 1839: 54 (*Clytus*)

disjunctus Pic, 1909: 123 (*Anaglyptus*)

Tribe **Clytini** Mulsant, 1839

Genus **Chlorophorus** Chevrolat, 1863: 290

type species *Callidium annulare* Fabricius, 1787

Subgenus **Chlorophoroides** Özdikmen, 2022: 637

type species *Leptura varia* O. F. Müller, 1766

damascenus Chevrolat, 1854: 483 (*Clytus*)

Sakarya, Balıkesir, Bursa, Tekirdağ

varius O. F. Müller, 1766: 188 (*Leptura*)
verbasci Linné, 1767 (*Leptura*)
nigrofasciatus Goeze, 1777: 507 (*Leptura*)
ornatus Herbst, 1784: 98 (*Callidium*)
gammoides Geoffroy, 1785: 81 (*Leptura*)
c-duplex Scopoli, 1786: 46 (*Stenocorus*)
venustus Gmelin, 1790: 1856 (*Cerambyx*)
strigosus Gmelin, 1790: 1877 (*Leptura*)
ferrugineus Mulsant, 1839: 87 (*Clytus*)
viridicollis Kraatz, 1871a: 410 (*Clytus*)
mixtornatus Fleischer, 1908: 211 (*Clytanthus*)
fontanae Hubenthal, 1923: 141 (*Clytanthus*)
incanus Plavilstshikov, 1924: 229
mouinieri Pic, 1950c: 6

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
Edirne, Kırklareli,
İstanbul (Thrace)

Subgenus **Crassofasciatus** Özdikmen, 2011a: 538
type species *Callidium trifasciatum* Fabricius, 1781

aegyptiacus Fabricius, 1775: 194 (*Callidium*) #
nigripes Brullé, 1832: 255 (*Clytus*)
magdalena Théry, 1895: cclxvi (*Clytus*)
rufifrons Pic, 1917a: 5
macedonicus Pic, 1935: 15

İstanbul (Anatolia), Balıkesir,
Bursa, Çanakkale (Anatolia),
İstanbul (Thrace)

convexifrons Holzschuh, 1981: 100 # İstanbul (Anatolia), Çanakkale (Anatolia)

hungaricus Seidlitz, 1891a: 828 Kocaeli, Bursa

niehuisi Adlbauer, 1992: 497 # *** Bilecik, Bursa

robustior Pic, 1900a: 11 (*Clytanthus*) # *** Bilecik, Bursa

trifasciatus Fabricius, 1781: 244 (*Callidium*)
portugallus Gmelin, 1790: 1854 (*Cerambyx*)
balearicus Pic, 1908a: 91 (*Clytanthus*)
perfidus Breit, 1915: 355 [DA]
paradoxus Dayrem, 1924: 78 (*Clytanthus*)

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik

Subgenus **Humeromaculatus** Özdikmen, 2011a: 536
type species *Cerambyx figuratus* Scopoli, 1763

dominici Sama, 1996a: 110 # *** İstanbul (Anatolia)

figuratus Scopoli, 1763: 55 (*Cerambyx*)
rusticus O. F. Müller, 1776: 93 (*Leptura*) [HN]
lamda Schrank, 1776: 67 (*Leptura*)
plebejus Fabricius, 1781: 243 (*Callidium*)
funebri Laicharting, 1784: 111 (*Clytus*)
leucozonias Gmelin, 1790: 1846 (*Cerambyx*)
cordiger Aragona, 1830: 26 (*Clytus*)
latifasciatus Fischer von Waldheim, 1832: 439 (*Clytus*)
conglobatus Fügner, 1891: 201 (*Clytus*)

Kocaeli, İstanbul (Anatolia), Kırklareli

tapaensis Pic, 1924: 22 (*Clytanthus*)

Subgenus **Perderomaculatus** Özdikmen, 2011a: 537
type species *Cerambyx sartor* Müller, 1766

sartor O. F. Müller, 1766: 188 (*Cerambyx*)
massiliensis Linnaeus, 1767: 1067 (*Leptura*)
rusticus Geoffroy, 1785: 81 (*Leptura*)
lineola Scopoli, 1786: 44 (*Stenocorus*)
achilleae Brahm, 1790: 141 (*Leptura*)
angusticollis Mulsant, 1851a: 123 (*Clytus*)
fulvicollis Mulsant, 1862: 181 (*Clytus*)
spinosulus Mulsant, 1862: 182 (*Clytus*)
griseus Gabriel, 1910: 556 (*Clytanthus*)
ruficeps Pic, 1917a: 5
mequignoni Pic, 1945b: 13

İstanbul (Anatolia), Balıkesir,
Bilecik, Bursa, Çanakkale
(Anatolia), Yalova,
Tekirdağ, Kırklareli,
İstanbul (Thrace)

Subgenus **Viridiphorus** Zamoroka, 2021: 207
type species *Callidium herbstii* Brahm, 1790
Brevenotatus Özdikmen, 2022: 635 type species *Clytus distinguendus* Perroud, 1855 (India)

herbstii Brahm, 1790: 148 (*Leptura*)
verbasci Fabricius, 1775: 194 (*Callidium*) [HN]
sulphureus Schaum, 1862: 103 (*Clytus*) [RN]

İstanbul (Anatolia), Balıkesir,
Çanakkale (Anatolia),
İstanbul (Thrace)

Genus **Clytus** Laicharting, 1784: 88
type species *Leptura arietis* Linnaeus, 1758

Subgenus **Clytoides** Özdikmen, 2023: 16
type species *Clytus rhamni* Germar, 1817

rhamni temesiensis Germar, 1824: 519 (*Callidium*)
ferruginipes Pic, 1891c: 26
longicollis Reitter, 1904: 82

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
Yalova, Tekirdağ, Kırklareli,
İstanbul (Thrace)

Subgenus **Clytus** Laicharting, 1784: 88
type species *Leptura arietis* Linnaeus, 1758

Europa J. Thomson, 1861: 221 type species *Leptura arietis* Linnaeus, 1758
Clytummus J. Thomson, 1861: 404 [RN] type species *Leptura arietis* Linnaeus, 1758
Sphegesthes Chevrolat, 1863: 333 type species *Leptura arietis* Linnaeus, 1758

arietis arietis Linnaeus, 1758: 399 (*Leptura*)
arcuatus Sulzer, 1761: 12 (*Leptura*)
quadrifasciatus DeGeer, 1775: 81 (*Cerambyx*)
gazella Fabricius, 1793: 333 (*Callidium*)
bourdilloni Mulsant, 1839: 81
cloueti Théry, 1893: ccxiii
bickhardti Pic, 1913: 98
heyrowskyi Pic, 1931b: 14

Kocaeli, İstanbul (Anatolia),
Bilecik, Çanakkale (Anatolia),
Kırklareli, İstanbul (Thrace)

chapmani Pic, 1937: 13

Subgenus ***Insolitomaculatus*** Özdikmen, 2023: 26
type species *Callidium tropicus* Panzer, 1795

tropicus Panzer, 1795: 265 (*Callidium*)
mucronatus Laporte & Gory, 1841: 52
interruptus Pic, 1931b: 14
prescutellaris Pic, 1931b: 14

Edirne

Genus ***Echinocerus*** Mulsant, 1862: 143
type species *Cerambyx floralis* Pallas, 1773
Paraplagionotus Kasatkin, 2005: 51 [unnecessary RN]

aulicus Laicharting 1784: 103 (*Clytus*)
basicornis Reitter, 1890b: 213 (*Plagionotus*)
controversus Schrank, 1798: 679 (*Clytus*)
massiliensis Pic, 1951: 1 (*Plagionotus*)

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
Yalova, Edirne, Kırklareli,
İstanbul (Thrace)

Genus ***Isotomus*** Mulsant, 1862: 183
type species *Callidium semipunctatum* Fabricius, 1798 (= *Callidium speciosum* Schneider, 1787)
Caloclytus Fairmaire, 1864a: 145 [RN] type species *Callidium semipunctatum* Fabricius, 1798
(= *Callidium speciosum* D. H. Schneider, 1787)

comptus comptus Mannerheim, 1825: 36 (*Clytus*)
perspicillus Fischer von Waldheim, 1832: 438 (*Clytus*)
pubicollis Laporte & Gory, 1841: 73 (*Clytus*)
insularis Laporte & Gory, 1841: 74 (*Clytus*)

Balıkesir

speciosus speciosus Schneider, 1787: 125 (*Callidium*)
semipunctatus Fabricius, 1798: 151 (*Callidium*)
stierlinii Tournier, 1872: 276 (*Clytus*)

Sakarya

Genus ***Neoplagionotus*** Kasatkin, 2005: 51
type species *Clytus bobelayei* Brullé, 1832

bobelayei bobelayei Brullé, 1832: 253 (*Clytus*)

İstanbul (Thrace)

bobelayei huseyini Lazarev, 2016a: 13
speciosus Adams, 1817: 309 (*Callidium*) [HN]

İstanbul (Anatolia)

Genus ***Plagionotus*** Mulsant, 1842a: 1
type species *Leptura detrita* Linnaeus, 1758
Platynotus Mulsant, 1839: 71 [HN] type species *Leptura detrita* Linnaeus, 1758

arcuatus arcuatus Linnaeus, 1758: 399 (*Leptura*)
lunatus Fabricius, 1782: 500 (*Callidium*)
salicis Schrank, 1798: 677 (*Clytus*)
reichei J. Thomson, 1861: 220 (*Plagyonotus*)

İstanbul (Thrace)

apicalis Hampe, 1863: 289 (*Clytus*)
stauropolibus Pic, 1915b: 7
martialis Pic, 1918: 15
buyssoni Dauphin, 1924: 42
pagnioni Pic, 1925a: 10
milliati Pic, 1934a: 20
plavilstshikovi Schmidt, 1951: 15
interrupteconnatus Schmidt, 1951: 16

arcuatus tastani Özdikmen, Atak & Uçkan, 2017: 89 # ***

Kocaeli, İstanbul
(Anatolia), Balıkesir,
Bilecik, Çanakkale
(Anatolia)

detritus detritus Linnaeus, 1758: 399 (*Leptura*)

convertini L. Petagna, 1819: 38 (*Callidium*)
rufescens Pic, 1891c: 24
obscrebasalis Pic, 1942a: 2
anticereductus Schmidt, 1951: 14
apicebimaculatus Schmidt, 1951: 14
uralensis Tippmann, 1952a: 144

Kocaeli, İstanbul (Anatolia),
İstanbul (Thrace)

Genus ***Xylotrechus*** Chevrolat, 1860b: 456
type species *Clytus sartorii* Chevrolat, 1860

Subgenus ***Rusticoclytus*** Vives, 1977: 130
type species *Leptura rustica* Linnaeus, 1758

rusticus Linnaeus, 1758: 398 (*Leptura*)

licii Linnaeus, 1767: 636 (*Cerambyx*)
hafniensis Fabricius, 1775: 190 (*Callidium*)
villosiflavomaculatus Goze, 1777: 507 (*Leptura*)
oo Schrank, 1781a: 162 (*Leptura*)
confusus Herbst, 1784: 98 (*Callidium*)
variegatus Geoffroy, 1785: 79 (*Leptura*)
signatus Geoffroy, 1785: 81 (*Leptura*)
longipes Villers, 1789: 256 (*Cerambyx*)
omega Rossi, 1790: 154 (*Callidium*)
octonotatus Gmelin, 1790: 1846 (*Cerambyx*)
maculatus Gmelin, 1790: 1877 (*Leptura*)
atomarius Fabricius, 1793: 332 (*Callidium*)
heros Ganglbauer, 1882: 138 (*Clytus*)
subuniformis Pic, 1902b: 15
basinotatus Pic, 1934a: 20
brevetestaceus Pic, 1934a: 20
fauconneti Pic, 1934a: 20
viturati Pic, 1934a: 20
apiceocellatus Schmidt, 1951: 13

Kocaeli, Sakarya, İstanbul
(Anatolia), Balıkesir,
Bursa, Yalova, Kırklareli,
İstanbul (Thrace)

Subgenus ***Xylotrechus*** Chevrolat, 1860b: 456
type species *Clytus sartorii* Chevrolat, 1860

Fulvotrechus Zamoroka, 2021: 205 type species *Xylotrechus stebbingi* Gahan, 1906

Hieroglyphotrechus Zamoroka, 2021: 205 type species *Clytus antilope* Schönherr, 1817
Ignetrechus Zamoroka, 2021: 205 type species *Xylotrechus pyrrhoderus* Bates, 1873
Spinotrechus Zamoroka, 2021: 204 type species *Clytus grayii* White, 1855

- antilope antilope** Schoenherr, 1817: 465 (*Clytus*) İstanbul (Anatolia), Bahkesir,
hieroglyphicus Drapiez, 1819: 294 (*Callidium*) [HN] Bursa, Kırklareli,
sekerai Paulian, 1986: 95 İstanbul (Thrace)
lento Paulian, 1986: 96
- arvicola arvicola** Olivier, 1800: 64 (*Callidium*) Bursa, Tekirdağ, Kırklareli
heydeni Stierlin, 1864: 152 (*Clytus*)
kraatz Lederer, 1864: 485 (*Clytus*)
tridentatus Bleuse, 1905: 21 (*Clytus*)
inbasalis Pic, 1911b: 6
herzegovinus Pic, 1913: 98
subangulosus Pic, 1934a: 21§
abbreviatus Schmidt, 1951: 14

Subfamily STENOPTERINAE Gistel, 1848

Tribe Stenopterini Gistel, 1848

Genus **Callimoxys** Kraatz, 1863: 105
type species *Stenopterus gracilis* Brullé, 1832

gracilis Brullé, 1832: 257 (*Stenopterus*) Tekirdağ

Genus **Callimus** Mulsant, 1846: [5]
type species *Callimus bourdini* Mulsant, 1846 (= *Saperda angulata* Schrank, 1789)

Subgenus **Callimus** Mulsant, 1846: [5]
type species *Callimus bourdini* Mulsant, 1846 (= *Saperda angulata* Schrank, 1789)
Callimellum Strand, 1928: 2 [unnecessary substitute name]

angulatus angulatus Schrank, 1789: 77 (*Saperda*) İstanbul (Anatolia), Bilecik,
cyaneus Fabricius, 1793: 330 (*Callidium*) Kırklareli, İstanbul (Thrace)
variabilis Bonelli, 1812: 181 (*Necydalis*)
laetus Motschulsky, 1845: 85 (*Stenopterus*)
bourdini Mulsant, 1846: [7]
gallicus Pic, 1932: 25 (*Callimellum*)

Subgenus **Lampropterus** Mulsant, 1862: 214
type species *Necydalis femoratus* Germar, 1823
Callimomimus İeniştea, 1952: 719 type species *Callimomimus distinctus* İeniştea, 1952 (= *Necydalis femoratus* Germar, 1823)

femoratus Germar, 1824: 519 (*Necydalis*) İstanbul (Anatolia), Bahkesir,
adonis Abeille de Perrin, 1881: 139 Bursa, Çanakkale (Anatolia),
narcissus Abeille de Perrin, 1881: 138 Edirne, Kırklareli,
thoracicus Chevrolat, 1882: 57 (*Callimoxys*) İstanbul (Thrace)
rumelicus Apfelbeck, 1899: 292

distinctus İeniştea, 1952: 719 (*Callimomimus*)

Genus **Stenopterus** Illiger, 1804: 120
type species *Necydalis rufa* Linnaeus, 1767

Subgenus **Bicallosocollus** Özdikmen, 2025i: 2766
type species *Stenopterus flavicornis* Küster, 1846

flavicornis Küster, 1846: 75
procerus Costa, 1855: 64

Edirne

kraatzii Pic, 1892n: 21 # ***

Balıkesir, Çanakkale (Anatolia)

Subgenus **Stenopterus** Illiger, 1804: 120
type species *Necydalis rufa* Linnaeus, 1767

rufus geniculatus Kraatz, 1863: 104

Kocaeli, İstanbul (Anatolia), Bilecik,
Bursa, Çanakkale (Anatolia), Yalova,
İstanbul (Thrace)

rufus rufus Linnaeus, 1767: 642 (*Necydalis*)
attenuatus Geoffroy, 1785: 84 (*Leptura*) [HN]

Kırklareli, İstanbul (Thrace)

Tribe **Psebiini** Lacordaire, 1868

Genus **Nathrius** Bréthes, 1916: 76
type species *Nathrius porteri* Bréthes, 1916 (= *Leptidea brevipennis* Mulsant, 1839)
Leptidea Mulsant, 1839: 105 [HN] type species *Leptidea brevipennis* Mulsant, 1839
Leptideella Strand, 1936: 169 [unnecessary substitute name]
Deuteroleptidea Paclt, 1946: 169 [unnecessary substitute name]

brevipennis Mulsant, 1839: 105 (*Leptidea*)
minutus Motschulsky, 1845: 84 (*Leptidea*)
manca LeConte, 1851: 24 (*Gracilia*)
rufipennis Dufour, 1851: 351 (*Gracilia*)
porteri Bréthes, 1916: 77

İstanbul (Anatolia), İstanbul (Thrace)

Tribe **Molorchini** Gistel, 1848

Genus **Molorchus** Fabricius, 1793: 356
type species *Necydalis umbellatarum* Schreber, 1759

Subgenus **Caenoptera** C. G. Thomson, 1859b: 150
type species *Necydalis minor* Linnaeus, 1758

minor minor Linnaeus, 1758: 421 (*Necydalis*)
ceramboides DeGeer, 1775: 151 (*Necydalis*)
dimidiatus Fabricius, 1775: 199 (*Leptura*)
medius Schrank, 1798: 688 (*Gymnopterion*)
rufescens Kiesenwetter, 1879: 316 [1879b: 60]

Bilecik, Bursa

Subgenus **Molorchus** Fabricius, 1793: 356

type species *Necydalis umbellatarum* Schreber, 1759

Glaphyra Newman, 1840a: 19 type species *Glaphyra semiusta* Newman, 1840

Laphyra Newman, 1842b: 418 [RN] type species *Glaphyra semiusta* Newman, 1840

Linomius Mulsant, 1862: 226 type species *Necydalis umbellatarum* Schreber, 1759

Sinolus Mulsant, 1862: 228 type species *Molorchus kiesenwetteri* Mulsant & Rey, 1861b

Conchopteris Fairmaire, 1864a: 153 type species *Necydalis umbellatarum* Schreber, 1759

Epanioglaphyra Niisato, 1986: 304 type species *Glaphyra kurosawai* Niisato, 1986

kiesenwetteri hircus Abeille de Perrin, 1881: 133

Bilecik, Bursa, Edirne

angorensis Pic, 1912d: 3 (*Caenoptera*)

anatolica Adlbauer, 1988: 275

umbellatarum umbellatarum Schreber, 1759: 9 (*Necydalis*)

Edirne

minimus Scopoli, 1763: 57 (*Necydalis*)

depressus Motschulsky, 1845: 85

diversipes Pic, 1897e: 31

Tribe **Dolocerini** Özdikmen, 2016

Genus **Dolocerus** Mulsant, 1862: 230

type species *Dolocerus reichii* Mulsant, 1862

Brachypteroma Heyden, 1863: 128 type species *Brachypteroma ottomanum* Heyden, 1863

reichii Mulsant, 1862: 231

mulsanti Stierlin, 1866: 30 (*Molorchus*)

ottomanus Heyden, 1863: 128 (*Brachypteroma*)

İstanbul (Anatolia), Balıkesir,
Bursa, İstanbul (Thrace)

Subfamily **LAMIINAE** Latreille, 1825

Tribe **Parmenini** Mulsant, 1839

Genus **Parmena** Dejean, 1821: 108

type species *Lamia unifasciata* Rossi, 1790

Subgenus **Parmena** Dejean, 1821: 108

type species *Lamia unifasciata* Rossi, 1790

istanbulensis Danilevsky & Hizal, 2017: 2 # ***

İstanbul (Thrace)

Tribe **Mesosini** Mulsant, 1839

Genus **Mesosa** Latreille, 1829: 124

type species *Cerambyx curculionoides* Linnaeus, 1760

Subgenus **Aplocnemina** Stephens, 1831: 236

type species *Cerambyx nubilus* Gmelin, 1790 (= *Lamia nebulosa* Fabricius, 1781)

Aphelocnemina Stephens, 1832: [414] [emendation, not in usage]

Haplocnemina Agassiz, 1846: 172 [unjustified emendation]

Helixoea Pascoe, 1865: 124 type species *Agelasta rupta* Pascoe, 1862

nebulosa nebulosa Fabricius, 1781: 218 (*Lamia*)

m-nigrum Schaller, 1783: 291 (*Cerambyx*)

brevis Villers, 1789: 257 (*Cerambyx*)

nubila Gmelin, 1790: 1832 (*Cerambyx*)

nebulator Turton, 1806: 306 (*Cerambyx*)

lineatopunctata Baudi di Selve, 1889: 193

silvestrii Tippmann, 1956: 492

İstanbul (Anatolia), Kırklareli,
İstanbul (Thrace)

Subgenus **Mesosa** Latreille, 1829: 124

type species *Cerambyx curculionoides* Linnaeus, 1760

Dendrobis Gistel, 1834: 30 [unnecessary substitute name]

curculionoides Linnaeus, 1760: 193 (*Cerambyx*)

oculata Geoffroy, 1785: 78 (*Leptura*) [HN]

tokatensis Pic, 1904b: 6 (*Haplocnemia*)

nigronotata Pic, 1906a: 86 (*Haplocnemia*)

Kırklareli

Tribe **Phrynetini** J. Thomson, 1864

Genus **Phrynet** Dejean, 1835: 341

type species *Lamia marmorea* Olivier, 1797

Inesida J. Thomson, 1860: 86 type species *Lamia leprosa* Fabricius, 1775

leprosa Fabricius, 1775: 178 (*Lamia*)

brunicornis Guérin-Ménéville, 1844: 239 (*Lamia*)

Kocaeli

Tribe **Batocerini** J. Thomson, 1864

Genus **Batocera** Dejean, 1835: 341

type species *Cerambyx rubus* Linnaeus, 1758

Megacriodes Pascoe, 1866: 259, 271 type species *Megacriodes saundersii* Pascoe, 1866

Semibatocera Kriesche, 1915: 115 type species *Batocera calana*, Parry, 1844 (= *Lamia calana* Parry, 1844)

Tyrannolamia Kriesche, 1915: 115 type species *Batocera wallacei* Thomson, 1858

rufomaculata DeGeer, 1775: 107 (*Cerambyx*)

cruentata Gmelin, 1790: 1863 (*Cerambyx*)

chlorinda J. Thomson, 1857a: 171

thysbe J. Thomson, 1878: 52

diana Nonfried, 1892a: 276

kuntzeni Kriesche, 1915: 139

flavescens Gilmour & Dibb, 1948: 68

Bursa

Tribe **Monochamini** Gistel, 1848

Genus **Anoplophora** Hope, 1839: 43

type species *Anoplophora stanleyana* Hope, 1839

Oplophora Hope, 1839: 42 type species *Oplophora sollii* Hope, 1839

Callophophora J. Thomson, 1864: 76 [unnecessary replacement name]

Cyriocrates J. Thomson, 1868: 181 type species *Oplophora horsfieldii* Hope, 1842

Melanauster J. Thomson, 1868: 181 type species *Cerambyx chinensis* Forster, 1771

Micromelanauster Pic, 1931c: 49 type species *Monochamus bowringii* White, 1858a
Falsocyriocrates Pic, 1953: 2 type species *Cyriocrates elegans* Gahan, 1888
Mimonemophas Breuning, 1961: 309 type species *Mimonemophas quadrifasciatus* Breuning, 1961

chinensis chinensis Forster, 1771: 39 (*Cerambyx*)
farinosa Houttuyn, 1766: 536 (*Cerambyx*) [HN]
punctator Fabricius, 1777: 230 (*Lamia*)
sinensis Gmelin, 1790: 1863 (*Cerambyx*)
abbreviata J. Thomson, 1865: 553 (*Callophophora*)
afflicta J. Thomson, 1865: 553 (*Callophophora*)
luctuosa J. Thomson, 1865: 553 (*Callophophora*)
sepulchralis J. Thomson, 1865: 553 (*Callophophora*)
perroudi Pic, 1953: 3 (*Melanauster*)

Sakarya, İstanbul (Anatolia),
İstanbul (Thrace)

glabripennis Motschulsky, 1854: 48 (*Cerosterna*)
laevigator J. Thomson, 1857c: 297 (*Cerosterna*)
nobilis Ganglbauer, 1889b: 82 (*Melanauster*)
angustata Pic, 1925b: 21 (*Melanauster*)
luteonotata Pic, 1925b: 21 (*Melanauster*)
nankinea Pic, 1926b: 2 (*Melanauster*)
laglaisei Pic, 1953: 3 (*Melanauster*)

İstanbul (Thrace)

malasiaca J. Thomson, 1865: 553 (*Callophophora*)

İstanbul (Anatolia)

Genus ***Monochamus*** Dejean, 1821: 106
type species *Cerambyx sutor* Linnaeus, 1758

Subgenus ***Monochamus*** Dejean, 1821: 106
type species *Cerambyx sutor* Linnaeus, 1758
Ceratades Gistel, 1834: 29 [unnecessary substitute name]
Monohammus Dejean, 1835: 340 [unjustified emendation]

galloprovincialis pistor Germar, 1818: 242 (*Lamia*)
lignator Krynicki, 1832: 158
nitidior Abeille de Perrin, 1870: 87 (*Monohammus*)
unifasciatus Pic, 1905b: 12 (*Monohammus*) ["Altař"]

İstanbul (Thrace)

galloprovincialis transitivus Lazarev, 2017: 18

Balıkesir, Bursa,
Çanakkale (Anatolia)

sutor sutor Linnaeus, 1758: 392 (*Cerambyx*)
atomarius DeGeer, 1775: 65 (*Cerambyx*)
heinrothi Cederhjelm, 1798: 88 (*Lamia*)
rosenmuelleri Cederhjelm, 1798: 89 (*Lamia*) [DA]
pellio Germar, 1818: 244 (*Lamia*)
obscurior Abeille de Perrin, 1869: 42 (*Monohammus*)

Balıkesir

Tribe **Lamiini** Latreille, 1825

Genus ***Herophila*** Mulsant, 1862: 273
type species *Dorcadion fairmairei* J. Thomson, 1857b

Dorcatypus J. Thomson, 1864: 77 type species *Dorcadion fairmairei* J. Thomson, 1857b

tristis obsoleta Fairmaire, 1859: 62 (*Morimus*) # *** İstanbul (Anatolia), Çanakkale (Anatolia), Kırklareli, İstanbul (Thrace), Çanakkale (Gelibolu)
interrupta Pic, 1921: 5

Genus ***Lamia*** Fabricius, 1775: 170

type species *Cerambyx textor* Linnaeus, 1758

Pachystola Dejean, 1833: 342 type species *Cerambyx textor* Linnaeus, 1758

Morimides J. Thomson, 1859a: 95 type species *Cerambyx textor* Linnaeus, 1758

textor Linnaeus, 1758: 392 (*Cerambyx*)

Sakarya, Edirne

noctis Gronov [Gronovius], 1764: 162 [n.538] (*Cerambyx*)

nigrorugosa DeGeer, 1775: 64 (*Cerambyx*)

unicolor P. Brown, 1776: 134 (*Cerambyx*) [HN]

tricarinata Cornelius, 1884: 50

Genus ***Morimus*** Brullé, 1832: 258

type species *Lamia lugubris* Fabricius, 1793 (= *Cerambyx asper* Sulzer, 1776)

Morinus Brullé, 1832: 258 [original incorrect spelling]

orientalis Reitter, 1894d: 43 #

Kocaeli, Sakarya, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa, Çanakkale
(Anatolia), Yalova, Kırklareli,
İstanbul (Thrace)

verecundus verecundus Faldermann, 1836: 396 (*Lamia*)

Balıkesir

ficitor Bogatchev, 1949: 183

Tribe **Pteropliini** J. Thomson, 1860

Genus ***Niphona*** Mulsant, 1839: 169

type species *Niphona picticornis* Mulsant, 1839

Subgenus ***Niphona*** Mulsant, 1839: 169

type species *Niphona picticornia* Mulsant, 1839

Aelara J. Thomson, 1864: 54 type species *Niphona regisfernandi* Paiva, 1860

Camptocnema J. Thomson, 1864: 54 type species *Niphona lateralis* White, 1858b

Ocheutes J. Thomson, 1864: 54 type species *Ocheutes scopulifera* J. Thomson, 1864

Falsoniphona Pic, 1925c: 26 type species *Falsoniphona lutea* Pic, 1925c

picticornis Mulsant, 1839: 169

Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia)

Tribe **Desmiphorini** J. Thomson, 1860

Genus ***Anaesthetis*** Dejean, 1835: 348

type species *Saperda testacea* Fabricius, 1781

Anaesthetus Gistel, 1856: 377 [unjustified emendation]

testacea rufescens Baeckmann, 1903: 395

Kocaeli, Sakarya, İstanbul
(Anatolia), Bursa

testacea testacea Fabricius, 1781: 235 (*Saperda*)

İstanbul (Thrace)

livida Herbst, 1784: 95 (*Saperda*)

fusca Geoffroy, 1785: 84 (*Leptura*)

teutonica Gmelin, 1790: 1842 (*Cerambyx*)

Tribe **Pogonocherini** Mulsant, 1839

Genus **Pogonocherus** Dejean, 1821: 107

type species *Cerambyx hispidus* Linnaeus, 1758

Subgenus **Pityphilus** Mulsant, 1862: 302

type species *Cerambyx ovatus* Goeze, 1777

decoratus Fairmaire, 1855: 320

İstanbul (Thrace)

Subgenus **Pogonocherus** Dejean, 1821: 107

type species *Cerambyx hispidus* Linnaeus, 1758

Pogonocherus Gistel, 1848b: 131 [unjustified emendation]

Strophinus Gistel, 1856: 376 type species *Cerambyx hispidus* Linnaeus, 1758

Pogonochaerus Gemminger, 1873: 3116 [unjustified emendation]

Eupogonocherus Linsley, 1935: 97 type species *Cerambyx hispidus* Linnaeus, 1758

eugeniae eugeniae Ganglbauer, 1891: 131 (*Pogonochaerus*)

pici Plavilstshikov, 1929: 34 (unavailable name).

Balıkesir,
İstanbul (Thrace)

hispidulus Piller & Mitterpacher, 1783: 35 (*Cerambyx*)

bidentatus C. G. Thomson, 1866: 85

bickhardti Pic, 1927b: 1 (*Pogonochaerus*)

İstanbul (Anatolia),
İstanbul (Thrace)

perroudi perroudi Mulsant, 1839: 158

vaulogerii Pic, 1927b: 1 (*Pogonochaerus*)

İstanbul (Anatolia), Balıkesir,
Bursa, Çanakkale (Anatolia),
Kırklareli, İstanbul (Thrace)

Tribe **Acanthoderini** J. Thomson, 1860

Genus **Aegomorphus** Haldeman, 1847: 45

type species *Aegomorphus decipiens* Haldeman, 1847 (= *Lamia modesta* Gyllenhal, 1817)

clavipes Schrank, 1781a: 135 (*Cerambyx*)

varius Fabricius, 1787: 130 (*Cerambyx*)

lucidus Starck, 1890: 71 (*Acanthoderes*)

Kocaeli, İstanbul (Anatolia), Bursa,
Edirne, İstanbul (Thrace)

Tribe **Acanthocinini** Blanchard, 1845

Genus **Acanthocinus** Dejean, 1821: 106

type species *Cerambyx aedilis* Linnaeus, 1758

Subgenus **Acanthocinus** Dejean, 1821: 106

type species *Cerambyx aedilis* Linnaeus, 1758

Aedilis Audinet-Serville, 1835a: 32 type species *Aedilis montanus* Audinet-Serville, 1835 (= *Cerambyx aedilis* Linnaeus, 1758)

Astynomus Dejean, 1835: 337 type species *Cerambyx aedilis* Linnaeus, 1758

Lamia Gistel, 1848b: xi [unjustified substitute name, HN] type species *Cerambyx aedilis* Linnaeus, 1758

Graphisurus Casey, 1913: 334 type species *Acanthocinus pusillus* Kirby, 1837

Canonura Casey, 1913: 335 type species *Aedilis spectabilis* LeConte, 1854

Tylocerina Casey, 1913: 335 type species *Cerambyx nodosus* Fabricius, 1775

Neacanthocinus Dillon, 1956: 230 type species *Cerambyx obsoletus* Olivier, 1795

aedilis Linnaeus, 1758: 392 (*Cerambyx*)

marmoratus Villers, 1789: 239 (*Cerambyx*)

montanus Audinet-Serville, 1835a: 33

obliteratus Pic, 1917a: 9

validus Matsushita, 1936: 148

dongbeiensis Z. Wang, 2003: 258

Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia),
İstanbul (Thrace)

griseus Fabricius, 1793: 261 (*Cerambyx*)

alpinus L.Redtenbacher, 1849: 494 (*Astynomus*)

obscurus Pic, 1891c: 32

novaki Tippmann, 1952b: 153

Kocaeli, Balıkesir, Bursa, Çanakkale
(Anatolia), Yalova, Edirne,
İstanbul (Thrace)

Genus **Leiopus** Audinet-Serville, 1835a: 86

type species *Cerambyx nebulosus* Linnaeus, 1758

Subgenus **Leiopus** Audinet-Serville, 1835a: 86

type species *Cerambyx nebulosus* Linnaeus, 1758

Liopus Agassiz, 1846: 212 [unjustified emendation]

Phrurus Gistel, 1856: 376 [unnecessary substitute name]

femoratus Fairmaire, 1859: 62 #

constellatus Mulsant & Rey, 1863: 159

caspius Ganglbauer, 1884: 531 (*Liopus*)

pachymerus Ganglbauer, 1884: 532 (*Liopus*)

Kocaeli, Sakarya, İstanbul (Anatolia),
Balıkesir, Çanakkale (Anatolia),
Yalova, Kırklareli, İstanbul (Thrace)

nebulosus caucasicus Ganglbauer, 1887a: 25 (*Liopus*) İstanbul (Anatolia), Balıkesir

nebulosus nebulosus Linnaeus, 1758: 391 (*Cerambyx*)

İstanbul (Thrace)

bifasciatus Goeze, 1777: 464 (*Cerambyx*) [HN]

monilis Geoffroy, 1785: 75 (*Cerambyx*)

fasciatus Villers, 1789: 239 (*Cerambyx*) [HN]

taeniatus Gmelin, 1790: 1863 (*Cerambyx*)

dissimilis Pic, 1889a: 5

unifasciatus Pic, 1891f: 23 (*Liopus*)

siculus Pic, 1924: 22 (*Liopus*)

Tribe **Exocentrini** Pascoe, 1864

Genus **Exocentrus** Dejean, 1835: 339

type species *Cerambyx balteatus* Fabricius *sensu* Dejean, 1835 (= *Cerambyx lusitanus* Linnaeus, 1767)

Oligopsis Thomson, 1864: 111 type species *Oligopsis exocentroides* J. Thomson, 1864
Camptomyne Pascoe, 1864b: 43 type species *Camptomyne callioides* Pascoe, 1864
Pseudocentrus Fairmaire, 1901: 230 type species *Exocentrus reticulatus* Fairmaire, 1898
Striatoexocentrus Breuning, 1955b: type species *Exocentrus nonymoides* Jordan, 1894
Formosexocentrus Breuning, 1958: 322 type species *Exocentrus variepennis* Schwarzer, 1925
Parasphigmothorax Breuning, 1974b: 155 type species *Parasphigmothorax ochreosignatus* Breuning, 1974
Bicolorihirtus Kusama & Tahira, 1978: 9 type species *Exocentrus venatoides* Kusama & Tahira, 1978

adspersus Mulsant, 1846: [9] [NP] Kocaeli, İstanbul (Anatolia), Balıkesir,
quercus Rossi, 1790: 143 (*Cerambyx*) Edirne, İstanbul (Thrace)
hirsutulus Faldermann, 1837: 272 (*Astynomus*) [DA]
clarae Mulsant & Rey, 1861a: 206 [1861c: 193]
revelierei Mulsant & Rey, 1875: 413 [=1875b: 77] (original spelling: *revelieri*)
alemdaghensis Breuning, 1981: 182

lusitanus Linnaeus, 1767: 1067 (*Cerambyx*) İstanbul (Anatolia), İstanbul (Thrace)
subpilosus Piller & Mitterpacher, 1783: 67 (*Cerambyx*)
pubicornis Schrank, 1789: 76 (*Cerambyx*)
crinitus Panzer, 1795: 249 (*Cerambyx*)
balteatus Gyllenhal, 1817: 163 (*Lamia*)
ochreoscutellatus Breuning, 1974b: 155

punctipennis punctipennis Mulsant & Guillebeau, 1856: 103 Edirne,
graecus Pic, 1901i: 52 İstanbul (Thrace)

punctipennis signatus Mulsant & Rey, 1863: 163 # İstanbul (Anatolia),
İstanbul (Thrace)

Tribe **Tetropini** Portevin, 1927

Genus **Tetrops** Kirby, 1826 (in Kirby & Spence 1826: 498)

type species *Leptura praeusta* Linnaeus, 1758

Subgenus **Tetrops** Kirby, 1826 (in Kirby & Spence 1826: 498)

type species *Leptura praeusta* Linnaeus, 1758

Anaetia Dejean, 1835: 350 type species *Leptura praeusta* Linnaeus, 1758

Polyopsia Mulsant, 1839: 190 type species *Leptura praeusta* Linnaeus, 1758

praeustus angorensis Pic, 1918: 11 # *** Sakarya, Bilecik

praeustus praeustus Linnaeus, 1758: 399 (*Leptura*) İstanbul (Anatolia),
pilosus Geoffroy, 1785: 78 (*Leptura*) [HN] İstanbul (Thrace)
ustulatus Hagenbach, 1822: 11 (*Saperda*)
inapicalis Pic, 1891c: 37
peterkai Scořepa, 2020: 565

praetermitus Sláma, 2020: 572

Tribe **Saperdini** Mulsant, 1839

Genus **Saperda** Fabricius, 1775: 184
type species *Cerambyx carcharias* Linnaeus, 1758

Subgenus **Compsidia** Mulsant, 1839: 182
type species *Cerambyx populneus* Linnaeus, 1758

populnea populnea Linnaeus, 1758: 394 (*Cerambyx*)
bilineata Scopoli, 1772: 102 (*Leptura*)
decempunctata DeGeer, 1775: 78 (*Cerambyx*)
betulina Geoffroy, 1785: 78 (*Leptura*)
salicis Zetterstedt, 1818: 258
populi Duméril, 1860: 607

Kocaeli, Sakarya, İstanbul
(Anatolia), Balıkesir, Bursa,
Çanakkale (Anatolia),
Edirne, İstanbul (Thrace)

quercus quercus Charpentier, 1825: 224

Edirne

Subgenus **Eusaperda** Danilevsky, 2023
type species *Cerambyx scalaris* Linnaeus, 1758

perforata Pallas, 1773: 723 (*Cerambyx*)
decempunctata Goeze, 1777: 506 (*Leptura*)
duodecimpunctata Brahm, 1790: 176 (*Cerambyx*)
seydlii Frölich, 1793: 135
rudolphi Cederhjelm, 1798: 92
algerica Pic, 1903b: 8
pallidipes Pic, 1904c: 9
albella Reitter, 1913b: 665

Kırklareli

scalaris scalaris Linnaeus, 1758: 394 (*Cerambyx*)
estellae Mulsant, 1839: 188
fenestrata Reineck, 1919: 72
algeriensis Breuning, 1952: 176
xantha Demelt, 1960: 182
niethammeri Podaný, 1963: 72

İstanbul (Thrace)

Subgenus **Lopezcolonia** Alonso-Zarazaga, 1998: 131 [RN]
type species *Saperda tremula* Fabricius, 1775 (= *Leptura octopunctata* Scopoli, 1772)
Argalia Mulsant, 1862: 381 [HN] type species *Saperda tremula* Fabricius, 1775 (= *Leptura octopunctata* Scopoli, 1772)
Nietzscheana Zubov, 2014: 695 type species *Nietzscheana plutenkoi* Zubov, 2014

octopunctata Scopoli, 1772: 101 (*Leptura*)
tremula Fabricius, 1775: 186
tiliae Schrank, 1798: 667
sempunctata Reitter, 1909: 57
magnini Dayrem, 1928a: 77

İstanbul (Thrace)

punctata Linnaeus, 1767: 1067 (*Cerambyx*)
gallica Pic, 1918: 5

İstanbul (Anatolia),
Kırklareli, İstanbul (Thrace)

Subgenus **Saperda** Fabricius, 1775: 184

type species *Cerambyx carcharias* Linnaeus, 1758

Anaerea Mulsant, 1839: 184 type species *Cerambyx carcharias* Linnaeus, 1758

Amilia Mulsant, 1862: 376 type species *Saperda phoca* Frölich, 1793 (= *Saperda similis* Laicharting, 1784)

carcharias Linnaeus, 1758: 394 (*Cerambyx*)
oppermanni Hummel, 1827: 13
villosa Gmelin, 1790: 1837 (*Cerambyx*)
grisescens Mulsant, 1839: 184 (*Anaerea*)

Bilecik, Bursa, İstanbul (Thrace)

Tribe **Phytoeciini** Mulsant, 1839

Genus **Oxylia** Mulsant, 1862: 398

type species *Phytoecia languida* Ménétriés, 1839

Coptosia Fairmaire, 1865: 177 type species *Phytoecia languida* Ménétriés, 1839

argentata argentata Ménétriés, 1832: 227 (*Saperda*)

Bursa

duponchelii Brullé, 1832: 260 (*Saperda*)

İstanbul (Thrace)

atomaria Frivaldszky von Frivald, 1837: 178 (*Saperda*)

vestita Küster, 1848a: 82 (*Phytoecia*)

Genus **Phytoecia** Dejean, 1835: 351

type species *Cerambyx cylindricus* Linnaeus, 1758

Subgenus **Helladia** Fairmaire, 1864a: 176

type species *Saperda flavescens* Brullé, 1832

humeralis humeralis Walzl, 1838: 471 (*Saperda*) #

frontalis Chevrolat, 1882: 63

scapularis Chevrolat, 1882: 63

Bursa, Edirne, Kırklareli,
İstanbul (Thrace)

millefolii Adams, 1817: 311 (*Saperda*)

azurea Steven, 1817: 190 (*Saperda*)

excelsa Stierlin, 1876: 479

Kocaeli, Bursa, Yalova,
Kırklareli, İstanbul (Thrace)

praetextata praetextata Steven, 1817: 184 (*Saperda*)

Kırklareli

Subgenus **Metacoptosia** Özdikmen, 2024: 3403

Type species *Phytoecia albovittigera* Heyden, 1863: 130

albovittigera Heyden, 1863: 130 #

Bilecik, İstanbul (Thrace)

languida Fairmaire, 1865: 177 (*Coptosia*) [HN]

vittigera Fairmaire, 1868: pl. 54, fig. 256 (*Coptosia*)

reichei Kraatz, 1876c: 287 (*Coptosia*)

semiannulicornis Pic, 1936: 4 (*Coptosia*)

Subgenus ***Metallicophytoecia*** Özdikmen, 2025e: 1819
type species *Leptura caerulea* Scopoli, 1772

baccueti Brullé, 1832: 262 (*Saperda*)

İstanbul (Anatolia), Balıkesir, Bilecik,
Bursa, Çanakkale (Anatolia), Yalova,
Edirne, Tekirdağ, İstanbul (Thrace)

caerulea Scopoli, 1772: 102 (*Leptura*)

İstanbul (Anatolia), Edirne,

viridis Gronov [Gronovius], 1764: 163 [n. 546] (*Cerambyx*)

Tekirdağ, İstanbul (Thrace)

rufimana Schrank, 1789: 77 (*Saperda*)

flavimana Creutzer, 1796: 15 (*Saperda*)

coelestis Townson, 1797: 469 (*Saperda*)

gilvimana Ménétriés, 1832: 227 (*Saperda*)

Subgenus ***Musaria*** J. Thomson, 1864: 120
type species *Leptura affinis* Herrer, 1784

affinis tuerki Ganglbauer, 1884: 575 #

Bursa, İstanbul (Thrace)

wachanrui Mulsant, 1851a: 127 #

Bilecik

syriaca Chevrolat, 1854: 484

alboscuteolata Chevrolat, 1854: 485

jezabel Reiche & Saulcy, 1858: 13

binodosa Chevrolat, 1882: 61

bisulcata Chevrolat, 1882: 61

tripunctata Chevrolat, 1882: 61

obscuricornis Pic, 1897c: 189

pallidithorax Pic, 1898g: 24 (*Musaria*)

rufomaculata Pic, 1917a: 9

jekeli Pic, 1917a: 10 (*Musaria*)

coyei Pic, 1952a: 695

berytensis Pic, 1952a: 696

breuningi Pic, 1952a: 696

haasi Pic, 1952a: 696

innotaticeps Pic, 1952a: 696 [HN]

irregularisignata Pic, 1952a: 696

marsabana Pic, 1952a: 696

microquadrisignata Pic, 1952a: 696

trinitatithorax Pic, 1952a: 696

Subgenus ***Neomusaria*** Plavilstshikov, 1928: 123
type species *Saperda balcanica* Frivaldszky von Frivald, 1835

balcanica balcanica Frivaldszky von Frivald, 1835: 268 (*Saperda*) İstanbul (Anatolia),
subvitticollis Breuning, 1951: 92 [Amasia] Bursa, İstanbul (Thrace)

Subgenus ***Neomusarioides*** Özdikmen, 2025c: 1800, 1801
type species *Saperda modesta* Waltl, 1838 = *Neomusaria waltli* Sama, 1991)

pauliraputii Sama, 1993: 295 (*Neomusaria*) # ***

Bilecik

waltli Sama, 1991: 127 [RN]

modesta Waltl, 1838: 471 (*Saperda*) [HN]

Bilecik

Subgenus **Opsilia** Mulsant, 1862: 387, 431

type species *Phytoecia flavicans* Mulsant, 1851 (= *Leptura coerulescens* Scopoli, 1763)

coerulescens coerulescens Scopoli, 1763: 49 (*Leptura*)

aeruginosa O. F. Müller, 1776: 95 (*Leptura*)

viridiuscula Goeze, 1777: 506 (*Leptura*)

virescens Fabricius, 1782: 499 (*Saperda*)

subcoerulea Geoffroy, 1785: 77 (*Leptura*)

trilineata Schoenherr, 1817: 433 (*Saperda*)

aeruginosa Mulsant, 1839: 210 [HN]

flavescens Mulsant, 1843: 284 [HN]

flavicans Mulsant, 1851a: 137 [RN]

grisescens Chevrolat, 1860c: 269

cobaltina Chevrolat, 1860c: 270

echii Chevrolat, 1860d: 302

chlorizans Chevrolat, 1860d: 303

incerta Mulsant, 1862: 433 (*Opsilia*)

obscura Brisout de Barneville, 1863: 116

nigrita Nedelkow, 1905: 204

estrelana Pic, 1930: 6 (*Opsilia*)

tienschanica Fuchs, 1965: 111

İstanbul (Anatolia),
Balıkesir, Bursa,
Çanakkale (Anatolia),
Tekirdağ, Kırklareli,
İstanbul (Thrace)

Subgenus **Paracoptosia** Danilevsky, 2017c: 1137

type species *Saperda compacta* Ménériés, 1832

bithynensis Ganglbauer, 1884: 573 #

trilinea Pic, 1892g: cxii

trilineata Pic, 1892g: cxii [= 1893d: 419]

conjuncta Pic, 1901c: 13 (*Coptosia*) [DA]

piciana Jakobson, 1924: 239 [RN]

amasina Pic, 1949: 60 (*Coptosia*)

limbata Pic, 1949: 61 (*Coptosia*) [DA]

latevittata Breuning, 1963b: 10 (*Conizonia*)

Bilecik, Bursa

Subgenus **Phytoecia** Dejean, 1835: 351

type species *Cerambyx cylindricus* Linnaeus, 1758

Hoplotoma Perez Arcas, 1874: 151 type species *Phytoecia malachitica* P. H. Lucas, 1847

cylindrica Linnaeus, 1758: 394 (*Cerambyx*)

cinerea DeGeer, 1775: 75 (*Cerambyx*)

verna O. F. Müller, 1776: 94 (*Leptura*)

silphoides Schrank, 1781a: 145 (*Cerambyx*)

fuliginosa Scopoli, 1786: 49 (*Leptura*)

simplonica Stierlin, 1878b: 438

grandis Pic, 1891h: 2 [DA]

kaemmereri Schatzmayr, 1928: 47

Kocaeli, İstanbul (Anatolia),
Kırklareli, İstanbul (Thrace)

geniculata geniculata Mulsant, 1862: 420 #

fuscicornis Mulsant & Rey, 1863: 168 [HN]

orientalis Kraatz, 1871a: 272 [RN]

İstanbul (Anatolia), Balıkesir,
Bilecik, Bursa,
Çanakkale (Anatolia)

donatellae Rapuzzi & Sama, 2010: 187

geniculata orientalis Kraatz, 1871c: 272 [RN] #

ingeniculata T. Pic, 1900b: 67

palaestina Pic, 1930: 3

Edirne, İstanbul (Thrace)

icterica Schaller, 1783: 292 (*Cerambyx*)

ephippium Fabricius, 1793: 317 (*Saperda*)

ragusana Küster, 1844: 55 (*Oberea*) [DA]

subannulipes Pic, 1910c: 51

İstanbul (Anatolia), Bilecik,
Tekirdağ, İstanbul (Thrace)

manicata Reiche & Saulcy, 1858: 17

Kocaeli, İstanbul (Anatolia),
Bursa, Çanakkale (Anatolia)

nigricornis Fabricius, 1782: 499 (*Saperda*)

melanocera Gmelin, 1790: 1838 (*Cerambyx*)

canaliculata Frölich, 1793: 144 (*Saperda*)

solidaginis Bach, 1856: 39

julii Mulsant, 1862: 429

caroni Mulsant & Godart, 1876: 419

tristriga Reitter, 1913a: 70

İstanbul (Thrace)

pubescens Pic, 1895d: 64

glaphyra K. Daniel, 1906: 177

İstanbul (Anatolia), Tekirdağ,
Kırklareli, İstanbul (Thrace)

pustulata pustulata Schrank, 1776: 66 (*Cerambyx*)

lineola Fabricius, 1781: 235 (*Saperda*)

posegana Piller & Mitterpacher, 1783: 67 (*Cerambyx*)

vulnerata Schaller, 1783: 293 (*Cerambyx*)

obscuripes Pic, 1895d: 65

macedonica Pic, 1929a: 9

brevenotata Pic, 1936: 4

Sakarya, İstanbul (Anatolia),
Bilecik, Edirne, Kırklareli

virgula virgula Charpentier, 1825: 225 (*Saperda*)

cyclops Küster, 1848b: 88

bravardi Pic, 1947: 1

İstanbul (Anatolia), Bilecik,
Bursa, Tekirdağ,
İstanbul (Thrace)

Genus ***Pilemia*** Fairmaire, 1864a: 175

type species *Phytoecia tigrina* Mulsant, 1851

Subgenus ***Pilemia*** Fairmaire, 1864a: 175

type species *Phytoecia tigrina* Mulsant, 1851

breverufonotata Pic, 1952c: 2 # ***

maculifera Holzschuh, 1984b: 170

Bursa

tigrina tigrina Mulsant, 1851a: 134 (*Phytoecia*)

anchusae Fuss, 1852: 138

Bilecik, Çanakkale (Anatolia)
European Turkey (? province)

Subgenus ***Pseudopilemia*** Kasatkin, 2018: 157

type species *Saperda hirsutula* Frölich, 1793

hirsutula hirsutula Frölich, 1793: 141 (*Saperda*)
atomaria Townson, 1797: 470 (*Saperda*)
tournieri Pic, 1952c: 2
androsensis Breuning, 1963b: 10 (*Oxyilia*)

Balıkesir, Bilecik, Kırklareli

Tribe **Obereini** Pascoe, 1864

Genus ***Oberea*** Dejean, 1835: 351
type species *Cerambyx linearis* Linnaeus, 1760

Subgenus ***Amaurostoma*** J. Müller, 1906: 223
type species *Cerambyx erythrocephalus* Schrank, 1776

erythrocephala erythrocephala Schrank, 1776: 67 (*Cerambyx*)
cincta Gebler, 1830: 186 (*Saperda*)
luteicollis Gebler, 1833: 303 (*Saperda*)
insidiosus Mulsant, 1862: 396
melitana Reiche, 1878: cxlix
richteri Bau, 1888: 425
anatolica Pic, 1901e: 19
hungarica Pic, 1914a: 11
montandoni Pic, 1914a: 11
theophilei Pic, 1914a: 11
erivanica Pic, 1917b: 11
planeti Pic, 1945a: 5
sinuatesignata Pic, 1945a: 7
schurmanni Heyrovský, 1962a: 42

Kocaeli, İstanbul
(Anatolia), Balıkesir,
Kırklareli

euphorbiae Germar, 1813: 131 (*Saperda*)
imitans G. Müller, 1948: 76
intermedia Breuning, 1947b: 59 [HN]

İstanbul (Thrace)

Subgenus ***Oberea*** Dejean, 1835 351
type species *Cerambyx linearis* Linnaeus, 1760
Isosceles Newman, 1842d: 318 type species *Isosceles macilentus* Newman, 1842

linearis Linnaeus, 1760: 191 (*Cerambyx*)
regularis Poda von Neuhaus, 1761: 38 (*Leptura*)
parallela Scopoli, 1763: 47 (*Leptura*)
fulvipes Geoffroy, 1785: 79 (*Leptura*)
cylindricollis Gray, 1832: 119 (*Saperda*)
limbata Mulsant, 1839: 197

İstanbul (Anatolia), İstanbul (Thrace)

oculata Linnaeus, 1758: 394 (*Cerambyx*)
inoculata Heyden, 1892: 81
borysthena Mokrzecki, 1900: 294
quadrinaculata Donisthorpe, 1913: 158
tomensis Kiseleva, 1927: 131

Kocaeli, Balıkesir, Kırklareli,
İstanbul (Thrace)

pupillata Gyllenhal, 1817: 185 (*Saperda*)
cognata Lentz, 1857: 145
alsatica Pic, 1915b: 10
luteonotata Pic, 1918: 5

Bilecik

Subfamily AGAPANTHIINAE Mulsant, 1839

Tribe Agapanthiini Mulsant, 1839

Genus **Agapanthia** Audinet-Serville, 1835a: 35
type species *Cerambyx cardui* Linnaeus, 1767

Subgenus **Agapanthia** Audinet-Serville, 1835a: 35
type species *Cerambyx cardui* Linnaeus, 1767

Eucrus Gistel, 1856: 376 type species *Cerambyx cardui* Linnaeus, 1767
Segmentaria Gistel, 1848b: viii [unnecessary substitute name]

cardui Linnaeus, 1767: 632 (*Cerambyx*)

marginalis Mulsant, 1839: 179
nigroaenea Mulsant, 1839: 179
consobrina Chevrolat, 1840: 17
peragalli Mulsant, 1862: 364
grossa Pic, 1891g: 63
pannonica Kratochvíl, 1985: 3

Kocaeli, İstanbul (Anatolia), Bursa,
Çanakkale (Anatolia), Edirne,
Kırklareli, İstanbul (Thrace)

suturalis Fabricius, 1787: 149 (*Saperda*)

annulata Fabricius, 1793: 314 (*Saperda*)
velox Gistel, 1857a: 560
subacutalis Chevrolat, 1882: 63
ruficornis Pic, 1918: 5
rufofemoralis Pic, 1946: 8 [Constantinople]

Kocaeli, Balıkesir, Bilecik, Bursa,
Çanakkale (Anatolia), Yalova,
Edirne, İstanbul (Thrace)

Subgenus **Epopetes** Gistel, 1857b: 93 [1857b: 605]

type species *Lamia asphodeli* Latreille, 1804

Agapanthiella Pesarini & Sabbadini, 2004b: 126 type species *Cerambyx villosoviridescens* DeGeer, 1775

asphodeli asphodeli Latreille, 1804: 282 (*Lamia*)

spencii Gyllenhal, 1817: 187 (*Saperda*)
reyi Mulsant & Godart, 1870: 27
insularis Gautier des Cottes, 1870: 263
delagrangei Pic, 1894e: 75 ["Akbès"]
mimica Pic, 1927b: 1

Balıkesir, Bilecik,
Çanakkale (Anatolia)

boeberi cynarae Germar, 1817: 222 (*Saperda*)

Kocaeli, İstanbul (Anatolia),
Balıkesir, Bilecik, Bursa,
Edirne, İstanbul (Thrace)

lateralis Ganglbauer, 1884: 541 # ***

bilateralis Pic, 1927b: 1
orientalis Pic, 1901h: 83 [DA]

Sakarya, İstanbul (Anatolia), Balıkesir,
Bilecik, Bursa, Çanakkale (Anatolia),
Edirne, Tekirdağ, İstanbul (Thrace)

villosoviridescens villosoviridescens DeGeer, 1775: 76 (*Cerambyx*) Sakarya, Balıkesir,
viridescens Gmelin, 1790: 1864 (*Cerambyx*) Bursa, Edirne, Tekirdağ
lineatocollis Donovan, 1797: 71 (*Cerambyx*)

latreillei ["latreille"] Fischer von Waldheim, 1805: 17 (*Saperda*)
angusticollis Gyllenhal, 1817: 189 (*Saperda*)
acutipennis Mulsant, 1862: 357
pyrenaee Brisout de Barneville, 1863: 117
nicaeensis Chevrolat, 1881: xcvi

Subgenus **Homoblephara** Pesarini & Sabbadini, 2004b: 128
type species *Saperda maculicornis* Gyllenhal, 1817

maculicornis maculicornis Gyllenhal, 1817, app.: 189 (*Saperda*) Balıkesir, Kırklareli
trivittata Gebler, 1830: 188 (*Saperda*)

Subgenus **Smaragdula** Pesarini & Sabbadini, 2004b: 128
type species *Saperda violacea* Fabricius, 1775

chalybea Faldermann, 1837: 303 İstanbul (Anatolia), Bilecik

frivaldszkyi Ganglbauer, 1884: 546 # Sakarya, İstanbul (Anatolia), Bilecik,
Bursa, İstanbul (Thrace)

osmanlis Reiche & Saulcy, 1858: 19 # İstanbul (Anatolia),
Bilecik, İstanbul (Thrace)

persicola Reitter, 1894c: 146 Sakarya
violaceipennis Pic, 1904c: 9

pesarinii Sama & Rapuzzi, 2010: 177 # Bilecik, Bursa, Çanakkale
(Anatolia), Tekirdağ

violacea Fabricius, 1775: 187 (*Saperda*) Kocaeli, Sakarya, İstanbul (Anatolia),
micans Fuessly, 1775: 13 (*Cerambyx*) Balıkesir, Bilecik, Bursa, Edirne,
cyanea Herbst, 1784: 95 (*Saperda*) Kırklareli, İstanbul (Thrace)
janthina Gmelin, 1790: 1842 (*Cerambyx*)
caerulea Schoenherr, 1817: 437 (*Saperda*)
smaragdina Krynicki, 1832: 161 (*Saperda*)
chalybaea Mulsant, 1839: 177 [HN]

Subgenus **Synthapsia** Pesarini & Sabbadini, 2004b: 121
type species *Saperda kirbyi* Gyllenhal, 1817

kirbyi zawadskyi Fairmaire, 1866b: 275 # Kocaeli, İstanbul (Anatolia), Bilecik,
valandovens Sláma, 2015c: 1127 Bursa, Çanakkale (Anatolia),
Edirne, İstanbul (Thrace)

Genus **Agapanthiola** Ganglbauer, 1900: 139
type species *Agapanthia euterpe* Ganglbauer, 1900 (= *Saperda leucaspis* Steven, 1817)
Calambiomorphus Pic, 1904b: 6 type species *Saperda leucaspis* Steven, 1817

leucaspis Steven, 1817: 184 (*Saperda*) İstanbul (Anatolia), Balıkesir,
cyarella Dalman, 1817b: 190 (*Saperda*) Edirne, İstanbul (Thrace)

pectoralis Eschscholtz, 1818: 482 (*Saperda*)
euterpe Ganglbauer, 1900: 139 (*Agapanthia*)

Genus **Calamobius** Guérin-Méneville, 1847: xviii
type species *Saperda gracilis* Creutzer, 1799 (= *Saperda filum* Rossi, 1790)

filum Rossi, 1790: 152 (*Saperda*)
hirtus Fabricius, 1793: 317 (*Saperda*)
gracilis Creutzer, 1799: 124 (*Saperda*)
marginellus Fabricius, 1802: 332 (*Saperda*)
tenuis Blanco Fernández, 1859: 411 (*Saperda*)
magnini Pic, 1931b: 10
decoloripes Pic, 1945a: 7

Kocaeli, İstanbul (Anatolia), Bilecik,
Bursa, Çanakkale (Anatolia),
İstanbul (Thrace)

Genus **Theophilea** Pic, 1895d: 39
type species *Theophilea cylindricollis* Pic, 1895

cylindricollis Pic, 1895d: 39 #
erzurumensis Önalp, 1974: 174 (*Agapanthia*)

Kocaeli

Subfamily **DORCADIONINAE** Swainson, 1840

Tribe **Dorcadionini** Swainson, 1840

Genus **Dorcadion** Dalman, 1817a: 397
type species *Cerambyx glycyrrhizae* Pallas, 1773

Subgenus **Carinatodorcadion** Breuning, 1943a: 524
type species *Cerambyx carinatus* Pallas, 1771

aethiops aethiops Scopoli, 1763: 53 (*Cerambyx*)
morio Fabricius, 1787: 141 (*Lamia*)
scopolii Gmelin, 1790: 1835 (*Cerambyx*)
bergmanni Tippmann, 1958: 151

Edirne

fulvum erythropterum Fischer von Waldheim, 1823: Tab. L - 7
canaliculatum Fischer von Waldheim, 1824: 240
frontale Mulsant & Rey, 1863: 154
opillicum Zamoroka, 2019: 365

İstanbul (Thrace)

hybridum hybridum Ganglbauer, 1884: 441

Edirne

ingae Peks, 1993: 3 # ***

Edirne, Tekirdağ, Kırklareli

Subgenus **Cribridorcadion** Pic, 1901c: 12

type species *Dorcadion mniszewi* Kraatz, 1873a

Pedestredorcadion Breuning, 1943a: 526 type species *Lamia pedestris* Poda von Neuhaus, 1761
Autodorcadion Plavilstshikov, 1958: 45 type species *Cerambyx arenarius* Scopoli, 1763
Dzhungarodorcadion Danilevsky, 1993: 47 type species *Dorcadion jacobsoni* Jakovlev, 1899a
Bergerianum Pesarini & Sabbadini, 2004c: 150 type species *Dorcadion chrysochroum* Breuning, 1943b

albolineatum Küster, 1847b: 86 # ***
byzantinum J. Thomson, 1867a: 69

Kocaeli, İstanbul (Anatolia),
Bilecik, İstanbul (Thrace)

anatolicum Pic, 1900b: 12 # ***
ininterruptum Pic, 1900b: 13 [HN]
mediointerruptum Pic, 1901d: 73
brevipenne Pic, 1902c: 10
cheshire Pic, 1902c: 10
subuniforme Pic, 1902d: 27

Çanakkale (Anatolia)

atritarse Pic, 1931b: 10

Çanakkale (Anatolia), Edirne

bangi heinzorum Braun, 1975: 17 # ***

Balıkesir

bithyniense Chevrolat, 1856a: 88 # ***
serotinum J. Thomson, 1865: 549

Sakarya, İstanbul (Anatolia),
Balıkesir, Bursa

breuningi Heyrovský, 1943: 78

Edirne

catenatum catenatum Waltl, 1838: 469 # ***
divisum Germar, 1839: 15
mancum Gistel, 1848: 431
subobliteratum T. Pic, 1899: 351
uninterruptum T. Pic, 1899: 351
dorsale T. Pic, 1907: 179
smyrnense Pic, 1917b: 10
bonyi Pic, 1942a: 1
smyrnanum Breuning, 1946: 106

Balıkesir, Bursa, Çanakkale (Anatolia)

catenatum dissimile Ganglbauer, 1884: 458 # ***

Çanakkale (Anatolia), Edirne,
Tekirdağ, Çanakkale (Gelibolu)

catenatum ezinense Özdiğmen & Skoupý, 2025b: 2045 # ***

Çanakkale (Anatolia)

catenatum loratum J. Thomson, 1867b: 123 # ***

Balıkesir, Çanakkale (Anatolia)

catenatum mytilenense Kraatz, 1873a: 66 #
latevittatum Kraatz, 1873a: 66
sparsedivisum Pic, 1911c: 185

Balıkesir

coiffaiti Breuning, 1962b: 392 # ***

Balıkesir

condensatum Küster, 1852: 92 # Kocaeli, İstanbul (Anatolia), İstanbul (Thrace)
punctatissimum Ganglbauer, 1884: 94
adampolum Pic, 1917a: 6
inlineatum Pic, 1917a: 6

completofascium Özdikmen & Skoupý, 2025a: 1996 # *** Çanakkale (Anatolia)

crux Billberg, 1817: 175 (*Lamia*) # *** İstanbul (Anatolia), Balıkesir, Bilecik, Bursa
smirnsensis Linnaeus, 1757: 411 (*Cerambyx*) [unavailable name]
smyrnsensis Linnaeus, 1762: 451 (*Cerambyx*) [unavailable name]
smyrnsensis Linnaeus, 1766: 229 (*Cerambyx*) [unavailable name]
graecum Waltl, 1838: 470
cruciator Gistel, 1848b: 131
bifidum Chevrolat, 1882: 60
integrofasciatum K. Daniel, 1900: 140
niveopictum K. Daniel, 1900: 140
lunulatum Pic, 1900b: 12
biinterruptum Pic, 1901e: 19
grandevittatum Pic, 1904c: 7
subfurcatum Pic, 1914a: 9
brevedisjunctum Pic, 1929a: 9
lateargenteum Pic, 1929a: 9

equestre bernhauerorum Peks, 2010: 213 # *** Bursa

equestre bisuturale Jureček, 1933: 128 Edirne

equestre reclinatum Kraatz, 1892: 173
immaculatum Kraatz, 1892: 174 ["Balkan"]

So far, I do not know any objective record from Turkey. However, it is reported by Danilevsky (2025) from Turkey in his catalogue, and so it can be distributed in European Turkey (e.g. Edirne, Kırklareli, Tekirdağ, İstanbul).

etruscum johannisfranci Pesarini & Sabbadini, 2007b: 40 Edirne

ferruginipes Ménétériés, 1836: 151 # Kocaeli, Sakarya, İstanbul (Anatolia),
thrasicum Küster, 1846: 90 İstanbul (Thrace)
sublineatum Pic, 1909: 99
subfuscopubens Pic, 1911b: 7

gallipolitanum gallipolitanum J. Thomson, 1867a: 59 # İstanbul (Anatolia),
suturalba Desbrochers des Loges, 1870: 125 Çanakkale (Anatolia),
gandolphei Tournier, 1872: 288 Edirne, Tekirdağ,
İstanbul (Thrace),
Çanakkale (Gelibolu)

gebzeense Breuning, 1974a: 148 # *** Kocaeli

infernale edremitense Breuning, 1966a: 20 # *** Balıkesir

- infernale infernale*** Mulsant & Rey, 1863: 158 # *** Bilecik, Kırklareli
rugosum J. Thomson, 1867a: 46 ["Siberia"]
revestitum K. Daniel, 1900: 140 ["Asia Minor"]
immutatum Pic, 1903c: 171 [Bos-Dagh, Eski-Chehir]
- inikliense*** Bernhauer & Peks, 2010b: 208 # *** Bursa
- intrusum*** Pesarini & Sabbadini, 2009: 26 # *** Edirne
- kindermanni*** Waltl, 1838: 470 # *** Çanakkale (Anatolia)
griseum Waltl, 1838: 470
- lineatocolle*** Kraatz, 1873a: 57 Edirne
atrofemoratum Pic, 1917a: 7
sinuaticolle Pic, 1927b: 6
riveti Pic, 1927a: 162
pseudolineatocolle Breuning, 1962b: 334
- lugubre*** Kraatz, 1873a: 41 Edirne
thessalicum Pic, 1916a: 22 ["Thessalie"]
pelionense Breit, 1923: 145 [Mt. Pelion]
discosulcatum Pic, 1926a: 1
sparsepilosum Pic, 1926a: 13 ["Salonique"]
pseudolugubre Breuning, 1943b: 96
minkovae Heyrovský, 1962b: 225 [Kresna]
meteorum Breuning, 1969b: 42 [Meteora]
parinfernale Breuning, 1975c: 10 [Kavala]
giachinoi Pesarini & Sabbadini, 2007b: 45 [Oros Kavala]
- micans subvestitum*** K. Daniel, 1900: 140 # *** Bursa, Çanakkale (Anatolia)
subreductum Pic, 1942a: 2
- obsoletum*** Kraatz, 1873a: 78 # İstanbul (Anatolia), Bursa, Edirne,
Tekirdağ, Kırklareli, İstanbul (Thrace)
- olympicum flavosuturale*** Krätschmer, 1987: 340 # *** Bursa
- olympicum olympicum*** Kraatz, 1873a: 78 # *** Bilecik, Bursa, Çanakkale
subalpinum Kraatz, 1873a: 78 (Anatolia), Kırklareli
oreophilum Ganglbauer, 1884: 500
- pedestre pedestre*** Poda von Neuhaus, 1761: 34 (*Cerambyx*) Balıkesir, Çanakkale
rufipes Fabricius, 1793: 288 (*Lamia*) (Anatolia), Kırklareli,
gogium J. Thomson, 1867b: 125 İstanbul (Thrace)
ruficornis Pic, 1931b: 10
zoufali Tippmann, 1958: 156
- pilosipenne kazdagense*** Özdikmen, 2013b: 56 # *** Balıkesir
- piochardi*** Kraatz, 1873a: 85 # *** Çanakkale (Anatolia)
quadripustulatum Ganglbauer, 1884: 457
elongatum Pic, 1898a: 58

pelops Jakovlev, 1901: 83
theophilei Pic, 1901g: 32 [HN]
roberti Pic, 1905b: 11
mauricei Plavilstshikov, 1927: 51 [RN]
varievittatum Pic, 1942c: 1
ladikense Breuning & Villiers, 1967a: 60 (*Pedestredorcadion*)

- pluto*** J. Thomson, 1867a: 47 # *** Sakarya
- praetermissum praetermissum*** Pesarini & Sabbadini, 1999: 47 # *** Bursa
- pseudobithyniense*** Breuning, 1962b: 465 # *** Balıkesir, Çanakkale (Anatolia)
- punctipenne*** Küster, 1852: 94 # Kocaeli, İstanbul (Anatolia), Bursa,
distinguendum Pic, 1931b: 10 ["Macédoine"] Çanakkale (Anatolia), İstanbul (Thrace)
- quadripustulatum*** Kraatz, 1873a: 88 # *** Balıkesir, Çanakkale (Anatolia)
- regulare regolare*** Pic, 1931b: 6 # Edirne, Kırklareli
- rolandmenradi*** Peks, 1992: 197 # *** Bursa
- scabricolle balikesirens*** Breuning, 1962b: 460 # *** Balıkesir, Bursa
- scabricolle crassofasciatum*** Özdikmen, 2013b: 56 # *** Balıkesir
- scabricolle uludaghicum*** Breuning, 1970: 98 # *** Bursa
- septemlineatum octolineatum*** Kraatz, 1873a: 61 # *** Kocaeli, Sakarya, İstanbul
(Anatolia), Balıkesir, Bilecik,
Bursa, Çanakkale (Anatolia),
Yalova
- septemlineatum septemlineatum*** Waltl, 1838: 469 # İstanbul (Thrace)
apicale J. Thomson, 1865: 549
rufopapicale Pic, 1942c: 2
- sinuatevittatum*** Pic, 1937: 6 # *** Balıkesir
brevemediolineatum Pic, 1942b: 6
- sturmii*** Frivaldszky von Frivald, 1837: 179 Edirne
- subinterruptum*** Pic, 1900b: 12 # *** Edirne
indivisum Pic, 1900b: 12
- tauricum pittinoi*** Pesarini & Sabbadini, 2010: 185 Edirne
- tauricum tauricum*** Waltl, 1838: 468 # Sakarya, İstanbul (Anatolia), Edirne,
nigritarse Kraatz, 1873a: 77 Tekirdağ, Kırklareli, İstanbul (Thrace)

- turcicum** Breuning, 1963a: 35 # *** Balıkesir
- Subgenus **Maculatodorcadion** Breuning, 1943a: 525
type species *Dorcadion quadrimaculatum* Küster, 1848a
- phrygicum** Peks, 1993: 8 # *** Balıkesir, Bursa, Çanakkale (Anatolia)
- quadrimaculatum nodicorne** Tournier, 1872: 287 # *** Çanakkale (Anatolia)
- quadrimaculatum quadrimaculatum** Küster, 1848a: 79 # Çanakkale (Gelibolu)
bimaculatum Pic, 1898a: 56
innotatum Pic, 1908d: 2
clementinum Pic, 1918: 11
- triste triste** Frivaldszky von Frivald, 1845: 184 # *** Balıkesir, Bursa
- wolffi** Krätschmer, 1985: 24 # *** Balıkesir
- Genus **Megalodorcadion** Pesarini & Sabbadini, 1999: 58
type species *Dorcadion ledereri* J. Thomson, 1865
- Subgenus **Anatolodorcadion** Özdikmen & Kaya, 2015: 3
type species *Dorcadion dombilicoides* Özdikmen & Kaya, 2013
- glabrofasciatum** K. Daniel, 1900: 140 # *** Balıkesir, Bilecik, Çanakkale (Anatolia)
imparivittatum K. Daniel, 1900: 140
nudefasciatum Özdikmen & Şenyüz, 2015: 647 (*Megalodorcadion*)
- Subgenus **Fusodorcadion** Özdikmen & Kaya, 2015: 3
type species *Dorcadion parallelum* Küster, 1847a
- parallelum** Küster, 1847a: 79 # *** İstanbul (Anatolia)
- Subgenus **Megalodorcadion** Pesarini & Sabbadini, 1999: 58
type species *Dorcadion ledereri* J. Thomson, 1865
- escherichi** Ganglbauer, 1897: 54 # *** Bilecik
egregium Ganglbauer, 1897: 56
angorensis Ganglbauer, 1897: 57
obliquesignatum Pic, 1900b: 12
postcedisjunctum Pic, 1909: 99
- Genus **Neodorcadion** Ganglbauer, 1884: 437
type species *Lamia bilineata* Germar, 1823
- Subgenus **Neodorcadion** Ganglbauer, 1884: 437
type species *Lamia bilineata* Germar, 1823
- bilineatum** Germar, 1824: 485 (*Lamia*)
macedonicum Pic, 1917a: 5 Edirne, Tekirdağ, Kırklareli, İstanbul (Thrace)

- difforme*** Pesarini & Sabbadini, 2014: 16 # *** Bursa
- exornatoides*** Breuning, 1962b: 68 # *** Balıkesir
- exornatum*** Frivaldszky von Frivald, 1835: 268 (*Dorcadion*) İstanbul (Anatolia),
labyrinthicum J. Thomson, 1865: 549 (*Dorcadion*) Balıkesir, Çanakkale
balcanicum Tournier, 1872: 284 (*Dorcadion*) (Anatolia), Edirne,
anticejunctum Pic, 1917a: 6 Kırklareli
subdiversum Pic, 1917a: 6
- fallax*** Kraatz, 1873a: 89 (*Dorcadion*) İstanbul (Anatolia), İstanbul (Thrace)
- laqueatum*** Walzl, 1838: 469 (*Dorcadion*) # Kocaeli, İstanbul (Anatolia),
interruptum Mulsant & Rey, 1863: 150 (*Dorcadion*) Balıkesir, Bursa, Çanakkale
sparsum Mulsant & Rey, 1863: 152 (*Dorcadion*) (Anatolia), Edirne,
abeillei Tournier, 1872: 208 (*Dorcadion*) İstanbul (Thrace)
anceyi Pic, 1913: 186
bulgaricum Pic, 1913: 186
diversesparsum Pic, 1914a: 8
insparsum Pic, 1914a: 8
lemoulti Pic, 1914a: 8
sparsenotatum Pic, 1914a: 8
servilorum Pic, 1917a: 6
subinterruptum Pic, 1917a: 6
kadiense Pic, 1917b: 10
- orientale*** Ganglbauer, 1884: 510 # *** Kocaeli, İstanbul (Anatolia)
flaschneri Pic, 1890a: clxxv
dispar Pic, 1892d: xxvii
merkli Pic, 1893b: cxcvi
eduardi Pic, 1900d: 82
- pelleti*** Mulsant & Rey, 1863: 149 (*Dorcadion*) İstanbul (Anatolia), İstanbul (Thrace)
segne Mulsant & Rey, 1863: 155 (*Dorcadion*)
disjunctum Pic, 1917a: 6
credulum Pic, 1917b: 10

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
Subgenus <i>Aegosoma</i>													
<i>scabricorne</i>			✓	✓	✓						✓	✓	
Tribe <i>Prionini</i>													
Genus <i>Mesoprionus</i>													
Subgenus <i>Trispinicollis</i>													
<i>besikanus</i>			✓		✓	✓	✓		✓			✓	
Genus <i>Prionus</i>													
<i>coriarius</i>	✓	✓	✓	✓	✓		✓				✓	✓	✓
Subfamily LEPTURINAE													
Tribe <i>Xylosteini</i>													
Genus <i>Xylosteus</i>													
<i>spinolae</i>											✓		
Tribe <i>Rhamnusiini</i>													
Genus <i>Rhamnusium</i>													
<i>bicolor graecum</i>												✓	
<i>bicolor juglandis</i>			✓			✓	✓						✓
Tribe <i>Evodinini</i>													
Genus <i>Brachyta</i>													
Subgenus <i>Fasciobrachyta</i>													
<i>balcanica</i>											✓		
Tribe <i>Acmaeopsini</i>													
Genus <i>Cortodera</i>													
<i>differens magdae</i>											✓		
<i>flavimana flavimana</i>	✓		✓							✓	✓	✓	
<i>flavimana inonuensis</i>						✓							
<i>humeralis humeralis</i>											✓		
<i>pumila meltemae</i>					✓						✓		
<i>rufipes rufipes</i>						✓							

[illegible]

[illegible]

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
Genus <i>Vadonia</i>													
Subgenus <i>Sinemaculipenna</i>													
<i>bitlisiensis</i>					✓								
Subgenus <i>Vadonia</i>													
<i>bolognai</i>	✓												
<i>grandicollis grandicollis</i>			✓									✓	
<i>moesiaca</i>			✓	✓					✓		✓		
<i>unipunctata unipunctata</i>			✓		✓				✓			✓	
Subfamily NECYDALINAE													
Tribe Necydalini													
Genus <i>Necydalis</i>													
Subgenus <i>Necydalis</i>													
<i>major</i>											✓		
<i>ulmi</i>											✓		
Subfamily SAPHANINAE													
Tribe Saphanini													
Genus <i>Saphanus</i>													
<i>kadleci</i>	✓	✓											
<i>piceus ganglbaueri</i>											✓		
Tribe Anisarthrini													
Genus <i>Alocerus</i>													
<i>moesiacus</i>				✓							✓		
Subfamily ASEMINEAE													
Tribe Asemini													
Genus <i>Arhopalus</i>													
<i>ferus</i>				✓	✓						✓		
<i>rusticus</i>			✓	✓			✓					✓	✓

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
Genus <i>Cephalocrius</i>													
<i>syriacus</i>				✓									
Subfamily SPONDYLIDINAE													
Tribe Spondylidini													
Genus <i>Spondylis</i>													
<i>buprestoides buprestoides</i>			✓	✓		✓						✓	
Subfamily CERAMBYCINAE													
Tribe Hesperophanini													
Subtribe Hesperophanina													
Genus <i>Hesperophanes</i>													
<i>sericeus</i>			✓	✓								✓	
Genus <i>Stromatium</i>													
<i>auratum</i>		✓	✓	✓		✓	✓		✓		✓	✓	
Genus <i>Trichoferus</i>													
Subgenus <i>Trichoferus</i>													
<i>fasciculatus fasciculatus</i>				✓		✓	✓						
<i>griseus</i>												✓	
<i>holosericeus</i>			✓									✓	
Tribe Phoracanthini													
Genus <i>Cordylomera</i>													
<i>spinicornis</i>	✓												
Tribe Cerambycini													
Genus <i>Cerambyx</i>													
Subgenus <i>Cerambyx</i>													
<i>cerdo acuminatus</i>	✓	✓	✓	✓	✓	✓	✓						
<i>cerdo cerdo</i>											✓	✓	
<i>dux</i>					✓	✓					✓	✓	

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
<i>miles</i>									✓			✓	
<i>nodulosus nodulosus</i>			✓			✓						✓	
<i>welensii welensii</i>		✓	✓	✓		✓						✓	
Subgenus <i>Microcerambyx</i>													
<i>scopolii scopolii</i>		✓	✓							✓	✓	✓	
Tribe <i>Compsocerini</i>													
Genus <i>Rosalia</i>													
Subgenus <i>Rosalia</i>													
<i>alpina</i>		✓	✓	✓			✓				✓	✓	
Tribe <i>Purpuricenini</i>													
Genus <i>Calchaenesthes</i>													
<i>oblongomaculata</i>												✓	
Genus <i>Purpuricenus</i>													
Subgenus <i>Basiatropansexus</i>													
<i>dalmatinus</i>					✓								
<i>inhumeralis</i>				✓			✓		✓				
<i>interscapillatus interscapillatus</i>						✓							
Subgenus <i>Purpuricenus</i>													
<i>budensis</i>	✓		✓	✓		✓	✓		✓			✓	
<i>kaehleri rossicus</i>			✓								✓	✓	
<i>neocaucasicus</i>						✓							
<i>renyvonaе latefasciatus</i>	✓		✓	✓		✓							
<i>renyvonaе renyvonaе</i>									✓		✓	✓	
<i>skypetarum zinaidae</i>	✓		✓										
Tribe <i>Callichromatini</i>													
Genus <i>Aromia</i>													
Subgenus <i>Aromia</i>													
<i>moschata moschata</i>	✓	✓	✓	✓		✓						✓	
<i>thoracica</i>				✓	✓	✓	✓					✓	

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
Subgenus <i>Phymatodes</i>													
<i>testaceus</i>			✓	✓		✓	✓		✓		✓	✓	
Subgenus <i>Poecilium</i>													
<i>alni alni</i>		✓	✓									✓	
Genus <i>Pyrrhidium</i>													
<i>sanguineum</i>												✓	
Genus <i>Ropalopus</i>													
Subgenus <i>Ropalopus</i>													
<i>clavipes</i>	✓		✓	✓	✓	✓		✓	✓		✓	✓	
<i>femoratus</i>										✓			
<i>macropus</i>	✓		✓								✓	✓	
<i>ungaricus insubricus</i>			✓									✓	
Genus <i>Semanotus</i>													
<i>russicus ruscicus</i>				✓									
Tribe <i>Anaglyptini</i>													
Genus <i>Anaglyptus</i>													
Subgenus <i>Anaglyptus</i>													
<i>arabicus</i>			✓										
<i>mysticus</i>											✓		
<i>obscurissimus</i>	✓												
Genus <i>Paraclytus</i>													
<i>sexguttatus</i>						✓					✓		
Tribe <i>Clytini</i>													
Genus <i>Chlorophorus</i>													
Subgenus <i>Chlorophoroides</i>													
<i>damascenus</i>		✓		✓		✓				✓			
<i>varius</i>	✓		✓	✓	✓	✓	✓		✓		✓	✓	
Subgenus <i>Crassofasciatus</i>													
<i>aegyptiacus</i>			✓	✓		✓	✓					✓	

[illegible]

[illegible]

[illegible]

[illegible]

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
Tribe <i>Saperdini</i>													
Genus <i>Saperda</i>													
Subgenus <i>Compsidia</i>													
<i>populnea populnea</i>	✓	✓	✓	✓		✓	✓		✓			✓	
<i>quercus quercus</i>									✓				
Subgenus <i>Eusaperda</i>													
<i>perforata</i>											✓		
<i>scalaris scalaris</i>												✓	
Subgenus <i>Lopezcolonia</i>													
<i>octopunctata</i>												✓	
<i>punctata</i>			✓								✓	✓	
Subgenus <i>Saperda</i>													
<i>carcharias</i>					✓	✓						✓	
Tribe <i>Phytoeciini</i>													
Genus <i>Oxylia</i>													
<i>argentata argentata</i>						✓							
<i>duponchelii</i>												✓	
Genus <i>Phytoecia</i>													
Subgenus <i>Helladia</i>													
<i>humeralis humeralis</i>						✓			✓		✓	✓	
<i>millefolii</i>	✓					✓		✓			✓	✓	
<i>praetextata praetextata</i>											✓		
Subgenus <i>Metacoptosia</i>													
<i>albovittigera</i>					✓							✓	
Subgenus <i>Metallicophytoecia</i>													
<i>baccueti</i>			✓	✓	✓	✓	✓	✓	✓	✓		✓	
<i>caerulea</i>			✓						✓	✓		✓	
Subgenus <i>Musaria</i>													
<i>affinis tuerki</i>						✓						✓	

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
<i>erythrocephala erythrocephala</i>	✓		✓	✓							✓		
<i>euphorbiae</i>												✓	
Subgenus <i>Oberea</i>													
<i>linearis</i>			✓									✓	
<i>oculata</i>	✓			✓							✓	✓	
<i>pupillata</i>					✓								
Subfamily AGAPANTHIINAE													
Tribe Agapanthiini													
Genus <i>Agapanthia</i>													
Subgenus <i>Agapanthia</i>													
<i>cardui</i>	✓		✓			✓	✓		✓		✓	✓	
<i>suturalis</i>	✓			✓	✓	✓	✓	✓	✓			✓	
Subgenus <i>Epoetes</i>													
<i>asphodeli asphodeli</i>				✓	✓		✓						
<i>boeberi cynarae</i>	✓		✓	✓	✓	✓			✓			✓	
<i>lateralis</i>		✓	✓	✓	✓	✓	✓		✓	✓		✓	
<i>villosoviridescens villosoviridescens</i>		✓		✓		✓			✓	✓			
Subgenus <i>Homoblephara</i>													
<i>maculicornis maculicornis</i>				✓							✓		
Subgenus <i>Smaragdula</i>													
<i>chalybea</i>			✓		✓								
<i>frivaldszkyi</i>		✓	✓		✓	✓						✓	
<i>osmanlis</i>			✓		✓							✓	
<i>persicola</i>		✓											
<i>pesarinii</i>					✓	✓	✓			✓			
<i>violacea</i>	✓	✓	✓	✓	✓	✓			✓		✓	✓	
Subgenus <i>Synthapsia</i>													
<i>kirbyi zawadskyi</i>	✓		✓		✓	✓	✓		✓			✓	

[illegible]

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
<i>completofascium</i>							✓						
<i>crux</i>			✓	✓	✓	✓							
<i>equestre bernhauerorum</i>						✓							
<i>equestre bisuturale</i>									✓				
<i>equestre reclinatum</i>													
<i>etruscum johannisfranci</i>									✓				
<i>ferruginipes</i>	✓	✓	✓									✓	
<i>gallipolitanum gallipolitanum</i>			✓				✓		✓	✓		✓	✓
<i>gebzeense</i>	✓												
<i>infernale edremitense</i>				✓									
<i>infernale infernale</i>					✓						✓		
<i>inikliense</i>						✓							
<i>intrusum</i>									✓				
<i>kindermanni</i>							✓						
<i>lineatocolle</i>									✓				
<i>lugubre</i>									✓				
<i>micans subvestitum</i>						✓	✓						
<i>obsoletum</i>			✓			✓			✓	✓	✓	✓	
<i>olympicum flavosuturale</i>						✓							
<i>olympicum olympicum</i>					✓	✓	✓				✓		
<i>pedestre pedestre</i>				✓			✓				✓	✓	
<i>pilosipenne kazdagense</i>				✓									
<i>piochardi</i>							✓						
<i>pluto</i>		✓											
<i>praetermissum praetermissum</i>						✓							
<i>pseudobithyniense</i>				✓			✓						
<i>punctipenne</i>	✓		✓			✓	✓					✓	
<i>quadripustulatum</i>				✓			✓						
<i>regulare regulare</i>									✓		✓		

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
<i>rolandmenradi</i>						✓							
<i>scabricolle balikesirense</i>				✓		✓							
<i>scabricolle crassofasciatum</i>				✓									
<i>scabricolle uludaghicum</i>						✓							
<i>septemlineatum octolineatum</i>	✓	✓	✓	✓	✓	✓	✓	✓					
<i>septemlineatum septemlineatum</i>												✓	
<i>sinuatevittatum</i>				✓									
<i>sturmii</i>									✓				
<i>subinterruptum</i>									✓				
<i>tauricum pittinoi</i>									✓				
<i>tauricum tauricum</i>		✓	✓						✓	✓	✓	✓	
<i>turcicum</i>				✓									
Subgenus <i>Maculatodorcadion</i>													
<i>phrygicum</i>				✓		✓	✓						
<i>quadrimaculatum nodicorne</i>							✓						
<i>quadrimaculatum quadrimaculatum</i>							✓						
<i>triste triste</i>				✓		✓							
<i>wolfi</i>				✓									
Genus <i>Megalodorcadion</i>													
Subgenus <i>Anatolodorcadion</i>													
<i>glabrofasciatum</i>				✓	✓		✓						
Subgenus <i>Fusodorcadion</i>													
<i>parallelum</i>			✓										
Subgenus <i>Megalodorcadion</i>													
<i>escherichi</i>					✓								
Genus <i>Neodorcadion</i>													
Subgenus <i>Neodorcadion</i>													
<i>bilineatum</i>									✓	✓	✓	✓	
<i>difforme</i>						✓							

	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
<i>exornatoides</i>				✓									
<i>exornatum</i>			✓	✓			✓		✓		✓		
<i>fallax</i>			✓									✓	
<i>laqueatum</i>	✓		✓	✓		✓	✓		✓			✓	
<i>orientale</i>	✓		✓										
<i>pelleti</i>			✓									✓	

	ASIAN TURKEY (NORTH-WEST ANATOLIA)								EUROPEAN TURKEY (THRACE)				
	KOCAELİ PART			SOUTH MARMARA PART					ERGENE PART		İSTRANCA PART	ÇATALCA PART	
	KOCAELİ	SAKARYA	İSTANBUL PARTLY	BALIKESİR	BİLECİK	BURSA	ÇANAKKALE PARTLY	YALOVA	EDİRNE	TEKİRDAĞ	KIRKLARELİ	İSTANBUL PARTLY	ÇANAKKALE PARTLY
TOTAL SPECIES-GROUP TAXA 332	70	38	136	111	72	113	84	22	66	25	94	150	8
MARMARA REGION	332												
KOCAELİ PART	161			-					-		-	-	
SOUTH MARMARA PART	-			205					-		-	-	
ERGENE PART	-			-					80		-	-	
İSTRANCA PART	-			-					-		94	-	
ÇATALCA PART	-			-					-		-	152	

With the present study, it is determined that Cerambycoidea taxa of Marmara region of Turkey includes a total of 332 species-group taxa of 93 subgenera of 98 genera belonging to 51 tribes of 11 subfamilies in 1 family as Cerambycidae [6 species-group taxa in Prioninae subfamily, 65 species-group taxa in Lepturinae subfamily, 2 species-group taxa in Necydalinae subfamily, 3 species-group taxa in Saphaninae subfamily, 3 species-group taxa in Aseminae subfamily, 1 species-group taxon in Spondylidinae subfamily, 76 species-group taxa in Cerambycinae subfamily, 12 species-group taxa in Stenopterinae subfamily, 73 species-group taxa in Lamiinae subfamily, 17 species-group taxa in Agapanthiinae subfamily, and 74 species-group taxa in Dorcadioninae subfamily]. Among 13 subfamilies of Cerambycoidea superfamily in Turkey, only Apatophyseinae in Cerambycidae family, and Vesperinae in Vesperidae family do not represent in longhorned beetles fauna of Marmara region of Turkey.

Accordingly, ratio of the number of species-group taxa of Marmara region of Turkey (332 taxa) to the number of total Cerambycoidea species-group taxa of Turkey (1073 taxa) is rather high and about 31 percent (30.94%).

Also, it is determined that Cerambycoidea taxa of Marmara region of Turkey includes a total of 70 endemic species-group taxa [7 endemic species-group taxa in Lepturinae subfamily, 1 endemic species-group taxon in Saphaninae subfamily, 9 endemic species-group taxa in Cerambycinae subfamily, 1 endemic species-group taxon in Stenopterinae subfamily, 5 endemic species-group taxa in Lamiinae subfamily, 1 endemic species-group taxon in Agapanthiinae subfamily, and 46 endemic species-group taxa in Dorcadioninae subfamily]. Among 11 subfamilies represented in this region of Cerambycoidea superfamily in Turkey, totally 4 subfamilies as Prioninae, Necydalinae, Aseminae, and Spondylidinae in Cerambycidae family do not include any endemic species-group taxon in longhorned beetles fauna of Marmara region of Turkey.

Accordingly, endemism ratio of Cerambycoidea taxa of Marmara region of Turkey is about 21 percent (21.08%). In addition, endemism ratio is 10.77 percent in Lepturinae subfamily, 33.33 percent in Saphaninae subfamily, 11.84 percent in Cerambycinae subfamily, 8.33 percent in Stenopterinae subfamily, 6.85 percent in Lamiinae subfamily, 5.88 percent in Agapanthiinae subfamily, and 62.16 percent in Dorcadioninae subfamily.

In addition, endemism ratio of the number of endemic species-group taxa of Marmara region of Turkey (70 taxa) to the number of total endemic Cerambycoidea species-group taxa of Turkey (536 taxa), and also to the total number of Cerambycoidea species-group taxa of Turkey (1073 taxa), are rather low, about 13 percent (13.06%) and about 7 percent (6.52%) respectively.

Moreover, it is determined that Cerambycoidea taxa of Marmara region of Turkey includes a total of 109 species-group taxa originally described from Turkey [14 species-group taxa in Lepturinae subfamily, 1 species-group taxon in Saphaninae subfamily, 1 species-group taxon in Aseminae subfamily, 14 species-group taxa in Cerambycinae subfamily, 1 species-group taxon in Stenopterinae subfamily, 15 species-group taxa in Lamiinae subfamily, 6 species-group taxa in Agapanthiinae subfamily, and 57 species-group taxa in Dorcadioninae subfamily]. Among 11 subfamilies represented in this region of Cerambycoidea superfamily in Turkey, totally 3 subfamilies as Prioninae, Necydalinae and Spondylidinae in Cerambycidae family do not include any species-group taxon originally described from Turkey in longhorned beetles fauna of Marmara region of Turkey.

Accordingly, ratio of species-group taxa originally described from Turkey (109 taxa) in Marmara region of Turkey to total Cerambycoidea species-group taxa originally described from Turkey (643 taxa) is low and about 17 percent (16.95%). In addition, about 64 percent (64.22%) of total species-group taxa originally described from Turkey in this region (109 taxa) is still endemic species-group taxa (70 taxa) to Turkey.

Longhorned beetles fauna of Asian Turkey (Anatolia) section of Marmara region of Turkey consists of 255 species-group taxa, and thereby about 77 percent (76.81%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in Asian Turkey (Anatolia) section.

Longhorned beetles fauna of Kocaeli Part of Asian Turkey (Anatolia) section of Marmara region of Turkey consists of 161 species-group taxa, and thereby about 49 percent (48.49%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in Kocaeli Part.

Longhorned beetles fauna of South Marmara Part of Asian Turkey (Anatolia) section of Marmara region of Turkey consists of 205 species-group taxa, and thereby about 62 percent (61.75%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in South Marmara Part.

Longhorned beetles fauna of European Turkey (Thrace) section of Marmara region of Turkey consists of 222 species-group taxa, and thereby about 67 percent (66.87%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in European Turkey (Thrace) section.

Longhorned beetles fauna of Ergene Part of European Turkey (Thrace) section of Marmara region of Turkey consists of 80 species-group taxa, and thereby about 24 percent (24.10%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in Ergene Part.

Longhorned beetles fauna of Istranca Part of European Turkey (Thrace) section of Marmara region of Turkey consists of 94 species-group taxa, and thereby about 28 percent (28.31%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in Istranca Part.

Longhorned beetles fauna of Çatalca Part of European Turkey (Thrace) section of Marmara region of Turkey consists of 152 species-group taxa, and thereby about 45 percent (45.18%) of total species-group taxa of Marmara region of Turkey (332 taxa) is represented in Çatalca Part.

The number of species-group taxa of the partial and provincial faunas of the parts and provinces in Marmara region of Turkey is as given at the end of the table above and the table below.

Longhorned beetles fauna of Marmara region of Turkey determined with the present study is lower numbers than expected. The main reason for this is that the region has not been studied sufficiently until now except for some provinces that have been studied sufficiently. It is clear that this region is still need to be investigated more comprehensively (as Çanakkale (Gelibolu), Yalova, Tekirdağ, Sakarya, Edirne, Kocaeli and Bilecik provinces especially).

According to Özdikmen (2008), the most recent study on longhorned beetles fauna of Marmara region of Turkey before this study, the fauna contained a total of 264 species-group taxa of 102 genera belonging to 7 subfamilies of 1 family as Cerambycidae. Whereas, according to the present study, a total of 332 species-group taxa of 98 genera belonging to 51 tribes of 11 subfamilies in 1 family as Cerambycidae are determined in longhorned beetles fauna of Marmara region.

Table 1. Surface areas of 11 provinces of 5 parts of 2 sections of Marmara region of Turkey researched in this study and their number of species-group taxa of Cerambycoidea.

PARTS AND PROVINCES	SURFACE AREA	NUMBER OF SPECIES-GROUP TAXA
SECTION ASIAN TURKEY (ANATOLIA)	49 484 km²	255
KOCAELİ PART	10 119 km²	161
KOCAELİ	3 397 km ²	70
SAKARYA	4 824 km ²	38
İSTANBUL partly	1 898 km ²	136
SOUTH MARMARA PART	39 365 km²	205
BALIKESİR	14 583 km ²	111
BİLECİK	4 179 km ²	72
BURSA	10 813 km ²	113
YALOVA	798 km ²	22
ÇANAKKALE partly	8 992 km ²	84
SECTION EUROPEAN TURKEY (THRACE)	23 182 km²	222
ERGENE PART	12 335 km²	80
EDİRNE	6 145 km ²	66
TEKİRDAĞ	6 190 km ²	25
İSTRANCA PART	6 459 km²	94

KIRKLARELİ	6 459 km ²	94
CATALCA PART	4 388 km ²	152
İSTANBUL partly	3 563 km ²	150
ÇANAKKALE partly (GELİBOLU)	825 km ²	8
MARMARA REGION	72 666 km ²	332

LITERATURE CITED

- Danilevsky, M. L. 2025. Catalogue of Palaearctic Chrysomeloidea (Vesperidae, Disteniidae, Cerambycidae). Retrieved October 30, 2025, from <https://www.cerambycidae.net/catalog.pdf>
- Della Beffa, G. 1915. I Coleotteri Italiani nocivi alle piante coltivate. Sistematica-Biologia-Agraria. Rivista Coleotterologica Italiana, 13 (1-3): appendix 1: 1-68.
- Özdikmen, H. 2021. An annotated catalogue: Cerambycoidea (Cerambycidae and Vesperidae) of Turkey (Coleoptera). Munis Entomology & Zoology, 16 (Supplement): 1273-1556.
- Özdikmen, H. 2024. Cerambycoidea of Turkey (Cerambycidae and Vesperidae) (Coleoptera). Munis Entomology & Zoology, 19 (suplement): 1153-2614.
- Özdikmen, H. 2025a. Host plants of Cerambycoidea of Turkey (Cerambycidae and Vesperidae) (Coleoptera). Munis Entomology & Zoology, 20 (1): 1-838.
- Özdikmen, H. 2025b. A taxonomic evaluation on the genus *Mesoprius* Jakovlev with a new subgenus (Cerambycidae: Prioninae). Munis Entomology & Zoology, 20 (2): 2998- 3002.
- Özdikmen, H. 2025c. A taxonomic evaluation on the genus *Vadonia* Mulsant with a new subgenus (Cerambycidae: Lepturinae). Munis Entomology & Zoology, 20 (2): 2982-2992.
- Özdikmen, H. 2025d. A taxonomic evaluation on *Spondylis buprestoides* (Linnaeus) (Coleoptera: Cerambycidae: Spondylidinae). Munis Entomology & Zoology, 20 (2): 2663- 2668.
- Özdikmen, H. 2025e. Proposition of a new subgenus of the genus *Trichoferus* Wollaston (Cerambycidae: Cerambycinae). Munis Entomology & Zoology, 20 (supplement): 4142- 4151.
- Özdikmen, H. 2025f. A taxonomic evaluation on the genus *Rosalia* Audinet-Serville (Cerambycidae: Cerambycinae). Munis Entomology & Zoology, 20 (2): 3303-3310.
- Özdikmen, H. 2025g. A taxonomic evaluation on the genus *Purpuricenus* Dejean with proposition of a new subgenus (Cerambycidae: Cerambycinae). Munis Entomology & Zoology, 20 (2): 2777-2798.
- Özdikmen, H. 2025h. Proposition of three new subgenera of the genus *Anaglyptus* Mulsant (Cerambycidae: Cerambycinae). Munis Entomology & Zoology, 20 (2): 2682-2709.

- Özdikmen, H. 2025i. A taxonomic evaluation on the genus *Stenopter* Illiger with proposition of two new subgenera (Cerambycidae: Stenopterinae). Munis Entomology & Zoology, 20 (2): 2761-2768.
- Özdikmen, H. 2025j. Taxonomic notes on Palaearctic longhorned beetles (Coleoptera: Cerambycidae). Munis Entomology & Zoology, 20 (supplement): 4002-4005.
- Özdikmen, H. 2026. A taxonomic evaluation on the genus *Aromia* Audinet-Serville with proposition of a new subgenus and a rearrangement on its species-group taxa (Cerambycidae: Cerambycinae). Munis Entomology & Zoology, 21 (2): 590-602.
- Özdikmen, H. & Özdikmen, G. 2025. A taxonomic evaluation on the genus *Aegosoma* Audinet-Serville with proposition of two new subgenera (Cerambycidae: Prioninae). Munis Entomology & Zoology, 20 (2): 2559-2580.
- Özdikmen, H. & Skoupý, V. 2025a. *Dorcadion* (*Cribridorcadion*) *completofascium* sp. nov. (Coleoptera: Cerambycidae: Dorcadioninae) from Turkey. Munis Entomology & Zoology, 20 (2): 1996-2003.
- Özdikmen, H. & Skoupý, V. 2025b. Two new subspecies of *Dorcadion catenatum* Waltl, 1838 (Coleoptera: Cerambycidae: Dorcadioninae) from Turkey. Munis Entomology & Zoology, 20 (2): 2041-2049.
- Zamoroka, A. M. 2022. Molecular revision of Rhagiini sensu lato (Coleoptera, Cerambycidae): Paraphyly, intricate evolution and novel taxonomy. Biosystems Diversity, 30 (3): 295–309.