Original article

doi:10.1111/codi.14373

Safety of primary anastomosis following emergency left sided colorectal resection: an international, multi-centre prospective audit

The 2017 European Society of Coloproctology (ESCP) collaborating group

European Society of Coloproctology (ESCP) Cohort Studies Committee, Hospital Universitari i Politècnic la Fe, València, Spain

Received 30 May 2018; accepted 6 August 2018

Abstract

Introduction Some evidence suggests that primary anastomosis following left sided colorectal resection in the emergency setting may be safe in selected patients, and confer favourable outcomes to permanent enterostomy. The aim of this study was to compare the major postoperative complication rate in patients undergoing end stoma *vs* primary anastomosis following emergency left sided colorectal resection.

Methods A pre-planned analysis of the European Society of Coloproctology 2017 audit. Adult patients (> 16 years) who underwent emergency (unplanned, within 24 h of hospital admission) left sided colonic or rectal resection were included. The primary endpoint was the 30-day major complication rate (Clavien-Dindo grade 3 to 5).

Results From 591 patients, 455 (77%) received an end stoma, 103 a primary anastomosis (17%) and 33 primary anastomosis with defunctioning stoma (6%). In multivariable models, anastomosis was associated with a similar major complication rate to end stoma (adjusted odds ratio for end stoma 1.52, 95%CI 0.83–2.79, $P = 0.173$). Although a defunctioning stoma was not associated with reduced anastomotic leak (12% defunctioned [4/33] *vs* 13% not defunctioned [13/97], adjusted odds ratio 2.19, 95%CI 0.43–11.02, $P = 0.343$), it was associated with less severe complications (75% [3/4] with defunctioning stoma, 86.7% anastomosis only [13/15]), a lower mortality rate (0% [0/4] *vs* 20% [3/15]), and fewer reoperations (50% [2/4] *vs* 73% [11/15]) when a leak did occur.

Conclusions Primary anastomosis in selected patients appears safe after left sided emergency colorectal resection. A defunctioning stoma might mitigate against risk of subsequent complications.

Keywords Surgery, emergency surgery, colon cancer, rectal cancer, gastrointestinal surgery, anastomotic leak, surgical complications, surgical outcomes

What does this paper add to the literature?
Anastomosis after emergency left sided colorectal resection is performed in up to one in five patients. In these highly selected patients, this study suggests that it is safe practice. A defunctioning stoma may mitigate against risk if an anastomotic leak subsequently occurs.

Introduction

In patients undergoing emergency left sided colorectal surgery, resection with end colostomy is a commonly described procedure. Concerns about the safety of any anastomosis in the emergency setting are particularly high in the presence of contamination or an unstable patient [1,2]. Although a stoma avoids the risk of anastomotic leak, it carries with its own morbidity and mortality profile (27–55% and 4–27% respectively) [2]. For patients that undergo end stoma formation, the reversal rate is as low as 44% in published series [3], with a significant impact on long-term quality of life and a risk of stoma-related complications.

Many studies have evaluated primary anastomosis in the emergency setting with generally favourable results. Multiple single-centre, retrospective, observational studies have demonstrated that anastomosis can be safely performed in selected patients within

Correspondence to: Luis Sánchez-Guillén, European Society of Coloproctology (ESCP) Cohort Studies Committee, Hospital Universitari i Politècnic la Fe, Avinguda de Fernando Abril Martorell, 106, 46026 València, Spain.
E-mail: drsanchezguillen@gmail.com

the emergency setting, even in presence of peritonitis [4–6]. However, the number of supporting randomised trials in the literature is low and those that exist are mainly related to peritonitis secondary to perforated diverticulitis, with primary anastomosis often only undertaken by specialised colorectal surgeons [4,7,8].

Decision-making about whether to create a primary anastomosis in selected, stable patients in an emergency setting remains a challenge for the individual surgeon. The decision must take into account patient comorbidities, intraoperative findings, underlying colorectal pathology, clinical status of the patient and expertise of the surgeon [9]. The aim of this multi-centre international study was to examine whether current decision-making in real-world settings supports primary anastomosis as a safe technique in selected patients after emergency left sided colorectal resection.

Methods

Protocol and centres

This prospective, observational, multi-centre study was conducted in line with a pre-specified protocol (http://www.escp.eu.com/research/cohort-studies). All participating centres were responsible for compliance to local approval requirements for ethics approval or indemnity as required. In the UK, the National Research Ethics Service tool recommended that this project was not classified as research, and the protocol was registered as clinical audit in all participating centres. Any unit performing gastrointestinal surgery was eligible to register to enter patients into the study. No minimum case volume, or centre-specific limitations were applied. The study protocol was disseminated to registered members of the European Society of Coloproctology (ESCP), and through national surgical or colorectal societies. This study represents planned analysis of the European Society of Coloproctology 2017 audit database.

Patient eligibility

Adult patients (> 16 years) undergoing left side colectomy or rectal resection, via any operative approach in emergency settings (within 24 h of hospital admission) were extracted from the ESCP 2017 Left Colon, Sigmoid and Rectal Resections Audit database. Any indication for surgery (benign or malignant) were eligible. Patients undergoing planned elective surgery were excluded, as were those undergoing left colorectal resection as part of a more extensive resection (e.g. subtotal colectomy, panproctocolectomy).




Figure 1 Flowchart for patients included in the analysis of postoperative outcomes of emergency colorectal surgery.

Data capture

Consecutive sampling was performed of eligible patients over an 8-week study period in each included centre. Local investigators commenced data collection on any date between the 1 January 2017 and 15 March 2017, with the last eligible patient being enrolled on 10 May 2017. Small teams of up to five surgeons or surgical trainees worked together to collect prospective data on all eligible patients at each centre. Quality assurance was provided by at least one consultant or attending-level surgeon. Data was recorded contemporaneously and stored on a secure, user-encrypted online platform (REDCap) without using patient identifiable information. Centres were asked to validate that all eligible patients during the study period had been entered, and to attain > 95% completeness of data field entry prior to final submission.

Outcome measure

The primary outcome measure was the 30-day postoperative major complication rate other, defined as Clavien-Dindo classification grade 3–5 (other than anastomotic leak including reoperation, reintervention, unplanned admission to critical care, organ support requirement or death). The secondary outcome measure was anastomotic leak, pre-defined as either (i)

Table 1 Patient and disease characteristics of included patients by anastomotic strategy.

Factor	Levels	Anastomosis, not defunctioned	Anastomosis, defunctioned	End stoma	*P*-value
Total number of patients		103	33	455	
Age group	< 55	26 (25.2)	11 (33.3)	72 (15.8)	0.056
	55–70	35 (34.0)	10 (30.3)	147 (32.3)	
	70–80	25 (24.3)	8 (24.2)	130 (28.6)	
	> 80	17 (16.5)	4 (12.1)	106 (23.3)	
Gender	Female	48 (46.6)	16 (48.5)	221 (48.6)	0.936
	Male	55 (53.4)	17 (51.5)	234 (51.4)	
ASA class	Low risk (ASA 1–2)	55 (53.4)	17 (51.5)	176 (38.7)	0.031
	High risk (ASA 3–5)	47 (45.6)	16 (48.5)	278 (61.1)	
BMI	Normal weight	31 (30.1)	14 (42.4)	135 (29.7)	0.353
	Underweight	1 (1.0)	1 (3.0)	16 (3.5)	
	Overweight	51 (49.5)	14 (42.4)	187 (41.1)	
	Obese	16 (15.5)	4 (12.1)	91 (20.0)	
History of IHD/CVA	No	86 (83.5)	33 (100.0)	363 (79.8)	0.013
	Yes	17 (16.5)	0 (0.0)	92 (20.2)	
History of diabetes mellitus	No	88 (85.4)	29 (87.9)	379 (83.3)	0.922
	Diabetes: any control	15 (14.6)	4 (12.1)	75 (16.5)	
Smoking history	Non-smoker	89 (86.4)	24 (72.7)	343 (75.4)	0.141
	Current	13 (12.6)	9 (27.3)	105 (23.1)	
Indication	Benign	62 (60.2)	26 (78.8)	325 (71.4)	0.042
	Malignant	41 (39.8)	7 (21.2)	130 (28.6)	
Resection type	Colonic only	67 (65.0)	16 (48.5)	271 (59.6)	0.315
	Involved rectum	35 (34.0)	17 (51.5)	183 (40.2)	
Approach	Laparoscopic	22 (21.4)	2 (6.1)	31 (6.8)	< 0.001
	Open	81 (78.6)	30 (90.9)	423 (93.0)	
	Robotic	0 (0.0)	1 (3.0)	0 (0.0)	
Training grade	Consultant	87 (84.5)	29 (87.9)	355 (78.0)	0.165
	Trainee	16 (15.5)	4 (12.1)	100 (22.0)	
Operator type	Colorectal	65 (63.1)	22 (66.7)	239 (52.5)	0.059
	General surgery	38 (36.9)	11 (33.3)	216 (47.5)	
Duration of surgery (minutes)	Mean (SD)	164.3 (73.3)	196.8 (58.2)	153.3 (63.6)	< 0.001

P-value derived from Chi-squared test for categorical variables and Student's t-tests for continuous variables after testing for normality. % shown by column. CVA, cerebrovascular accident; IHD, Ischemic heart disease; IQR, interquartile range; N/A, not applicable; SD, standard deviation. BMI groups are categorised as Underweight (< 18.5), Normal weight (18.5–25), Overweight (25–30), Obese (> 30).

gross anastomotic leakage proven radiologically or clinically, or (ii) the presence of an intraperitoneal (abdominal or pelvic) fluid collection on post-operative imaging.

Statistical analysis

This report has been prepared in accordance to guidelines set by the STROBE (strengthening the reporting of observational studies in epidemiology) statement for observational studies [10]. Patient, disease and operative characteristics were compared using Student's t-test for normal, continuous data, Mann–Whitney *U* test for non-normal continuous data or Chi-squared test for categorical data. To test the association between the outcome measures and the main explanatory variables of interest (expedited *vs* emergency, end stoma *vs* primary anastomosis), a mixed-effects logistic regression model was fitted. Clinically plausible patient, disease and operation-specific factors were entered into the model for risk-adjustment, treated as fixed effects. These were defined *a priori* within the study protocol, and included irrespective of their significance on univariate analysis. Hospitals were entered into the model as a random-effect, to adjust for hospital-level variation in outcome. Effect estimates are presented as odds ratios (OR) with 95% confidence intervals (95% CI) and two-tailed *P*-values. Model discrimination was quantified using *C*-statistic, or the area under the receiver operating characteristic curve (AUC) of the model. An alpha level of 0.05 was used throughout. Data analysis was undertaken using R Studio V3.1.1 (R Foundation, Boston, MA, USA).

Results

Patients

This study included 591 patients undergoing emergency surgery from 43 countries (Fig. 1). The mean age of patients was 67.4 years (ranging from 18 to 96). 51.8% were male and 57.4% had a high anaesthetic risk class (ASA 3–5). Differences in demographics between patients with anastomosis and end stoma are shown in Table 1. Primary anastomosis was performed in 136 patients (23%) with 33 of these patients receiving a defunctioning stoma. This stoma was a loop ileostomy in 84.8% (28/33), an end/double-barreled ileostomy in 6.1% (2/33) and a loop colostomy in 9.1% (3/33). 30.1% (178/591) of included operations were done for malignancy, with end stoma being most common operative strategy (73.0%, 130/178). Of these, 20.2% were undergoing neoadjuvant therapy prior to their presentation for emergency surgery (short course radiotherapy, 7/36; long course chemoradiotherapy, 18/36; chemotherapy only: 11/36). Primary anastomosis with or without defunctioning stoma was performed less frequently than end stoma in disease affecting the rectum (14.9% and 7.2% *vs* 77.9% respectively). An anastomosis was attempted in 27% (87/326) of patients operated upon by a colorectal surgeon and 18% (49/265) by a general surgeon ($P = 0.059$, Fig. 2).

Major complications (Clavien-Dindo 3 to 5)

Results of analysis for factors associated with the occurrence of major complications are shown in Table 2. An




Figure 2 Variation in anastomotic practice between colorectal and general surgeons.

Table 2 Univariable and multilevel models for major postoperative complications.

Factor	Levels	No major complication	Major complication	OR (univariable)	OR (multilevel)
Anastomosis type	Anastomosis, not defunctioned	79 (19.8)	18 (11.6)	– *(Reference)*	– *(Reference)*
	Anastomosis, defunctioned	28 (7.0)	5 (3.2)	0.78 (0.24–2.18, $P = 0.658$)	0.63 (0.20–2.02, $P = 0.442$)
	End stoma	292 (73.2)	132 (85.2)	1.98 (1.17–3.54, $P = 0.015$)	1.52 (0.83–2.79, $P = 0.173$)
Age	< 55	79 (19.8)	26 (16.8)	–	–
	55–70	135 (33.8)	45 (29.0)	1.01 (0.58–1.78, $P = 0.964$)	0.86 (0.46–1.60, $P = 0.635$)
	70–80	104 (26.1)	48 (31.0)	1.40 (0.81–2.48, $P = 0.236$)	1.03 (0.54–1.96, $P = 0.918$)
	> 80	81 (20.3)	36 (23.2)	1.35 (0.75–2.46, $P = 0.320$)	0.91 (0.45–1.82, $P = 0.784$)
Gender	Female	202 (50.6)	65 (41.9)	–	–
	Male	197 (49.4)	90 (58.1)	1.42 (0.98–2.07, $P = 0.067$)	1.66 (1.10–2.51, $P = 0.016$)
ASA class	Low risk (ASA 1–2)	192 (48.1)	40 (25.8)	–	–
	High risk (ASA 3–5)	207 (51.9)	115 (74.2)	2.67 (1.78–4.05, $P < 0.001$)	2.54 (1.59–4.07, $P < 0.001$)
BMI	Normal weight	119 (29.8)	60 (38.7)	–	–
	Underweight	11 (2.8)	7 (4.5)	1.26 (0.44–3.37, $P = 0.647$)	1.37 (0.46–4.07, $P = 0.566$)
	Overweight	194 (48.6)	55 (35.5)	0.56 (0.36–0.86, $P = 0.009$)	0.53 (0.33–0.85, $P = 0.009$)
	Obese	75 (18.8)	33 (21.3)	0.87 (0.52–1.45, $P = 0.603$)	0.76 (0.43–1.33, $P = 0.332$)
History of IHD/CVA	No	330 (82.7)	117 (75.5)	–	–
	Yes	69 (17.3)	38 (24.5)	1.55 (0.99–2.42, $P = 0.054$)	1.12 (0.67–1.87, $P = 0.669$)
History of diabetes mellitus	No	339 (85.0)	126 (81.3)	–	–
	Diabetes: any control	60 (15.0)	29 (18.7)	1.30 (0.79–2.10, $P = 0.292$)	0.99 (0.57–1.72, $P = 0.965$)
Smoking history	Non-smoker	311 (77.9)	120 (77.4)	–	–
	Current	88 (22.1)	35 (22.6)	1.03 (0.65–1.60, $P = 0.894$)	0.94 (0.57–1.55, $P = 0.802$)
Indication	Benign	276 (69.2)	114 (73.5)	–	–
	Malignant	123 (30.8)	41 (26.5)	0.81 (0.53–1.22, $P = 0.312$)	0.85 (0.54–1.34, $P = 0.481$)
Resection type	Colonic only	237 (59.4)	94 (60.6)	–	–
	Involved rectum	162 (40.6)	61 (39.4)	0.95 (0.65–1.38, $P = 0.788$)	1.01 (0.66–1.54, $P = 0.964$)
Approach	Open	354 (88.7)	148 (95.5)	–	–
	Minimally invasive	45 (11.3)	7 (4.5)	0.37 (0.15–0.79, $P = 0.018$)	0.42 (0.17–1.02, $P = 0.055$)
Training grade	Consultant	320 (80.2)	119 (76.8)	–	–
	Trainee	79 (19.8)	36 (23.2)	1.23 (0.78–1.90, $P = 0.373$)	1.01 (0.61–1.65, $P = 0.978$)
Operator type	Colorectal	218 (54.6)	87 (56.1)	–	–
	General surgery	181 (45.4)	68 (43.9)	0.94 (0.65–1.37, $P = 0.751$)	0.97 (0.62–1.51, $P = 0.888$)

Major postoperative complications were pre-defined as Clavien-Dindo grade complications 3 to 5 (re-operation, re-intervention, admission to critical care or death. Odds ratio (OR) presented with 95% confidence intervals. % shown by column. CVA, cerebrovascular accident; IHD, Ischemic heart disease; IQR, interquartile range; N/A, not applicable; SD, standard deviation.

end stoma was significantly associated with increased major postoperative complications upon univariable analysis (OR 1.98 95% CI 1.17–3.54, $P = 0.015$), but this association was not seen following risk adjustment (adjusted odds ratio for end stoma in mixed effects model 1.52, 95%CI 0.83–2.79, $P = 0.173$). In the multilevel model significant predictors for major complications were high ASA risk (grade 3–5) (OR 2.54, 95% CI 1.59–4.07, $P < 0.001$) and male gender (OR 1.66, 95% CI 1.10–2.51, $P = 0.016$). Overweight BMI was associated with a lower major complication rate than a normal BMI (OR 0.53, 95% CI 0.33–0.85, $P = 0.009$), however the location of resection (involving rectum or colonic only) demonstrated no association. The model demonstrated fair discrimination (AUC: 0.71).

Anastomotic leak

Unadjusted outcomes according the anastomotic strategy, stratified by presence of leak, are shown in Table 3. Although a defunctioning stoma was not associated with reduced anastomotic leak (12% defunctioned [4/33] *vs* 13% not defunctioned [13/97],

Table 3 Outcomes of patients undergoing emergency left sided colorectal surgery with or without anastomosis.

Factor	Levels	Anastomosis, defunctioned no leak	Anastomosis, defunctioned with leak	Anastomosis, no leak	Anastomosis, with leak	End stoma	*P*-value
Post-operative complication	No complication	15 (51.7)	0 (0.0)	50 (56.8)	0 (0.0)	159 (34.9)	< 0.001
	Minor complication (Clavien-Dindo 1–2)	12 (41.4)	1 (25.0)	30 (34.1)	2 (13.3)	149 (32.7)	
	Major complication (Clavien-Dindo 3–5)	2 (6.9)	3 (75.0)	8 (9.1)	13 (86.7)	147 (32.3)	
Post-operative mortality	No	28 (96.6)	4 (100.0)	88 (100.0)	12 (80.0)	390 (85.7)	0.001
	Yes	1 (3.4)	0 (0.0)	0 (0.0)	3 (20.0)	65 (14.3)	
Re-operation	No re-operation	28 (96.6)	2 (50.0)	83 (94.3)	4 (26.7)	405 (89.0)	< 0.001
	Re-operation	1 (3.4)	2 (50.0)	5 (5.7)	11 (73.3)	50 (11.0)	
Critical care admission	Missing	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	0 (0.0)	< 0.001
	None	15 (51.7)	2 (50.0)	63 (71.6)	6 (40.0)	179 (39.3)	
	Planned from theatre	13 (44.8)	2 (50.0)	19 (21.6)	9 (60.0)	216 (47.5)	
	Unplanned from theatre	1 (3.4)	0 (0.0)	5 (5.7)	0 (0.0)	49 (10.8)	
	Unplanned from ward	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	11 (2.4)	
Re-admission	No	28 (96.6)	3 (75.0)	78 (88.6)	14 (93.3)	422 (92.7)	0.746
	Yes	1 (3.4)	1 (25.0)	9 (10.2)	1 (6.7)	28 (6.2)	
	missing	0 (0.0)	0 (0.0)	1 (1.1)	0 (0.0)	5 (1.1)	
Length of stay	Mean (SD)	11 (6.2)	18.5 (9.1)	9 (4.3)	18.7 (6.4)	13.6 (7.8)	< 0.001

P-values derived from Chi-squared test for categorical variables and Student's *T*-test for parametric continuous variables, % shown by column.

adjusted odds ratio 2.19, 95%CI 0.43–11.02, $P = 0.343$), it was associated with fewer major complications (75% [3/4] with defunctioning stoma, 86.7% anastomosis only [13/15]), lower mortality (0% [0/4] *vs* 20% [3/15]), and reoperation 50% [2/4] *vs* 73% [11/15]) when a leak did occur (Fig. 3). The minor complication rate was similar between groups where the anastomosis successfully healed without leak (41.4% defunctioned [12/27] *vs* 34.1% not defunctioned [30/88]) and where an end stoma was formed (32.7% [149/455]). On the univariable analysis (Table 4) previous history of IHD/CVA (OR 5.06, 95% CI 1.50–16.27, $P = 0.007$) was associated with an increased risk of leak, whilst being of middle age was protective (age 55–70 years old; OR 0.10, 95% CI 1.61–100, $P = 0.037$). When a multilevel model was




Figure 3 Clavien Dindo complication grade, grouped by anastomotic outcome.

Table 4 Univariable and multilevel models for anastomotic leak amongst patients with anastomosis only.

Factor	Levels	No leak	Leak	OR (univariable)	OR (multilevel)
Defunctioning ileostomy	No	84 (74.3)	13 (76.5)	– *(Reference)*	– *(Reference)*
	Yes	29 (25.7)	4 (23.5)	0.89 (0.24–2.75, $p = 0.851$)	2.19 (0.43–11.02, $P = 0.343$)
Age	< 55	30 (26.5)	7 (41.2)	–	–
	55–70	42 (37.2)	1 (5.9)	0.10 (0.01–0.62, $P = 0.037$)	0.05 (0.00–0.66, $P = 0.023$)
	70–80	28 (24.8)	4 (23.5)	0.61 (0.15–2.25, $P = 0.470$)	0.32 (0.05–1.91, $P = 0.213$)
	> 80	13 (11.5)	5 (29.4)	1.65 (0.42–6.17, $P = 0.458$)	1.03 (0.15–6.96, $P = 0.980$)
Gender	Female	57 (50.4)	6 (35.3)	–	–
	Male	56 (49.6)	11 (64.7)	1.87 (0.66–5.74, $P = 0.249$)	1.50 (0.38–5.87, $P = 0.563$)
ASA class	Low risk (ASA 1–2)	62 (54.9)	7 (41.2)	–	–
	High risk (ASA 3–5)	51 (45.1)	10 (58.8)	1.74 (0.62–5.09, $P = 0.296$)	1.00 (0.20–4.96, $P = 0.996$)
History of IHD/CVA	No	102 (90.3)	11 (64.7)	–	–
	Yes	11 (9.7)	6 (35.3)	5.06 (1.50–16.27, $P = 0.007$)	5.10 (0.75–34.53, $P = 0.095$)
History of diabetes mellitus	No	99 (87.6)	12 (70.6)	–	–
	Diabetes: any control	14 (12.4)	5 (29.4)	2.95 (0.84–9.34, $P = 0.074$)	8.56 (1.16–63.38, $P = 0.035$)
Smoking history	Non-smoker	96 (85.0)	14 (82.4)	–	–
	Current	17 (15.0)	3 (17.6)	1.21 (0.26–4.21, $P = 0.782$)	1.44 (0.25–8.19, $P = 0.678$)
Indication	Benign	75 (66.4)	10 (58.8)	–	–
	Malignant	38 (33.6)	7 (41.2)	1.38 (0.47–3.89, $P = 0.543$)	1.26 (0.29–5.47, $P = 0.753$)
Resection type	Colonic only	64 (56.6)	14 (82.4)	–	–
	Involved rectum	49 (43.4)	3 (17.6)	0.28 (0.06–0.92, $P = 0.055$)	0.18 (0.03–1.00, $P = 0.050$)
Training grade	Consultant	98 (86.7)	12 (70.6)	–	–
	Trainee	15 (13.3)	5 (29.4)	2.72 (0.78–8.55, $P = 0.095$)	1.06 (0.19–5.95, $P = 0.944$)
Operator type	Colorectal	74 (65.5)	9 (52.9)	–	–
	General surgery	39 (34.5)	8 (47.1)	1.69 (0.59–4.75, $P = 0.319$)	2.19 (0.55–8.76, $P = 0.267$)

Overall anastomotic leak was pre-defined as either (i) gross anastomotic leakage proven radiologically or clinically, or (ii) the presence of an intraperitoneal (abdominal or pelvic) fluid collection on post-operative imaging. Odds ratio (OR) presented with 95% confidence intervals. % shown by column. SD, standard deviation; IQR, Interquartile range; IHD, Ischemic heart disease; CVA, cerebrovascular accident; N/A, not applicable.

fitted (Fig. 4), a history of diabetes also conveyed an increased risk of leak (OR 8.56, 95% CI 1.16–63.38, $P = 0.035$). The model demonstrated good discrimination (AUC: 0.87).

Discussion

This study showed that primary anastomosis was performed in up to one in five patients and appears safe in this highly selected group after emergency left sided colorectal resection (unplanned, within 24 h of hospital admission). A defunctioning stoma was only used in 24% of patients with a primary anastomosis. The exploratory findings of this study, limited by small numbers, suggested that a defunctioning stoma may mitigate against risk if an anastomotic leak occurs. Other patient-related risk characteristics (male gender, high ASA grade) and an open approach were identified as independent risk factors for major postoperative complications. Furthermore, young and elderly age or a history of diabetes were shown as risk factors for anastomotic leak in emergency procedures.

Previously, the simple formation of an end colostomy after resection of the pathology (ubiquitously known as a 'Hartmann's procedure') has been advocated as the gold standard treatment in emergency left colonic resection, to eliminate risk of anastomotic leak [11–13]. In the last 15 years, several studies have questioned this strategy [14,15]. A primary anastomosis is not only feasible, it may even be associated with better postoperative outcomes, both in terms of complications and mortality [16,17]. Given that more than 40% of temporary stomas become permanent, selecting patients correctly for a primary anastomosis is attractive [18–20]. In addition, reversal of Hartmann's can be a technically demanding operation resulting in further morbidity and mortality [21]. These findings support a recent consensus statements and prospective multi-centre randomized trials that suggest primary anastomosis with proximal diversion as an optimal strategy for sigmoid diverticulitis in selected patients with Hinchey 3 or 4 disease [4,22,23]. This current study gives credence to the current situation and confirms that surgeons are making appropriate decisions on a case-by-case level, thereby




Figure 4 Forest plot demonstrating mixed effects model for factors associated with anastomotic leak in patients undergoing emergency left sided colorectal surgery.

effectively stratifying patients for primary anastomosis or end stoma. It is known that a defunctioning stoma in elective surgery has utility in mitigating the clinical impact of anastomotic leak [24,25]. Loop ileostomies and their closure are not complication-free and several studies have shown that temporary loop ileostomies can become permanent in up to 25% of patients [24–28]. However this study comparably suggests that a defunctioning stoma may mitigate some risk when a leak occurs. This must be interpreted with caution since numbers in this study were low; for example, only four patients with an anastomosis and defunctioning stoma suffered a leak.

There were slightly more primary anastomotic attempts by colorectal *vs* general surgeons in this study. Even though there is no homogenous definition of colorectal surgeon internationally, results of multiple studies confirm the importance of colorectal specialisation in the emergency setting [29,30]. An individual surgeon's personality and their response to perceived operative risk may also influence choice of anastomotic strategy [31]. Further research is needed to determine whether the grade and surgical specialism of the operating (or senior) surgeon, and specialisation and experience of included centres affect both the decision for anastomosis and the subsequent clinical outcome.

There are inherent limitations to the 'snapshot' observational study reported here which we have attempted to overcome in the study design, statistical analysis and interpretation. There is an obvious selection bias in this study, although we planned the analysis around this *a priori*. We aimed to analyse safety of current practice; this study showed that end stoma was more frequently used in older patients, with poor general status, in smokers, and in those with arteriopathy and benign disease. However this paper defines outcomes in the highly selected group of patients undergoing anastomosis, and thus supports surgical decision making in specific cases, rather than in recommending a general change in approach. The low numbers of anastomotic leak and major complication within secondary analyses of the subgroup undergoing anastomosis (< 25% of included patients) makes estimation of effect sizes inaccurate here (reflected by broad confidence intervals). Therefore, this should be seen as exploratory only; the analysis would likely be underpowered to detect a small to moderate effect size. We are also unable to comment on the appropriateness of decision making and have not collected detailed information on parameters that may effect this (for example: contamination (Mannheim Peritonitis Index [32]), previous surgery, intraoperative physiological instability). Most of the literature available on this topic is based on retrospective or single centres data which lacks sufficient detail to allow case-mix adjustment in multivariable models. This study therefore adds to the literature in providing a contemporary perspective using a prospective international observational study design, with a pre-specified protocol and analysis plan. In addition, the variety of centers included (in terms of number of patients, facilities and different technologies available) in this study delivers a realistic picture of the current management of emergency left colorectal resections, reducing selection bias and increasing the external validity of the findings. The different countries and even continents involved ensured the result's validity resolving the demographic differences in diverticulitis and cancer across countries. Finally, the study is limited by short-term follow up to 30 days only; we have not collected data on stoma reversal rates, quality of life or stoma-related complications following surgery. An alternative complication categorisation system such as the Comprehensive Complications Index [33] may also give increased fidelity in comparisons between intermediate term outcomes. Further evaluation of these important parameters following emergency left sided colorectal surgery is warranted.

The data from this study supports current international practice of primary anastomosis following emergency left sided colorectal resection in a highly selected group of patients, demonstrating satisfactory safety and an acceptable morbidity profile. Where an anastomosis is formed, a defunctioning stoma does not appear to reduce the risk of leak, but may mitigate the severity of resultant complications.

Acknowledgements

Supported by the European Society of Coloproctology (ESCP). REDCap and infrastructural support was received from the Birmingham Surgical Trials Institute (BiSTC) at the Birmingham Clinical Trials Unit (BCTU).

Conflicts of interest

None to declare.

Funding

None.

References

1 Hartmann H. Note sur un procédé nouveau d'extirpation des cancers de la partie du côlon. *Bull Mem Soc Chir Paris* 1923; **49:** 1474–7.
2 Barbieux J, Plumerias F, Hammy A. Current indications for Hartmann procedure. *J Visc Surg* 2016; **153:** 31–8.
3 van de Wall BJ, Draaisma WA, Schouten ES, Broeders IA, Consten EC. Conventional and laparoscopic reversal of the Hartmann procedure: a review of literature. *J Gastrointest Surg* 2010; **14:** 743–52.
4 Bridoux V, Regimbeau JM, Ouaissi M *et al.* Hartmann's procedure or primary anastomosis for generalized peritonitis due to perforated diverticulitis: a prospective multicenter randomized trial (DIVERTI). *J Am Coll Surg* 2017; **225:** 798–805.
5 Salem L, Flum DR. Primary anastomosis or Hartmann's procedure for patients with diverticular peritonitis? A systematic review. *Dis Colon Rectum* 2004; **47:** 1953–64.
6 Trenti L, Biondo S, Golda T *et al.* Generalized peritonitis due to perforated diverticulitis: Hartmann's procedure or primary anastomosis? *Int J Colorectal Dis* 2011; **26:** 377–84.
7 Oberkofler CE, Rickenbacher A, Raptis DA *et al.* A multicenter randomized clinical trial of primary anastomosis or Hartmann's procedure for perforated left colonic diverticulitis with purulent or fecal peritonitis. *Ann Surg* 2012; **256:** 819–26; discussion 26–7.
8 Binda GA, Karas JR, Serventi A *et al.* Primary anastomosis vs nonrestorative resection for perforated diverticulitis with

peritonitis: a prematurely terminated randomized controlled trial. *Colorectal Dis* 2012; **14:** 1403–10.
9 Gibbons G, Tan CJ, Bartolo DC *et al.* Emergency left colonic resections on an acute surgical unit: does subspecialization improve outcomes? *ANZ J Surg* 2015; **85:** 739–43.
10 von Elm E, Altman DG, Egger M *et al.* Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007; **335:** 806–8.
11 Myers E, Hurley M, O'Sullivan GC, Kavanagh D, Wilson I, Winter DC. Laparoscopic peritoneal lavage for generalized peritonitis due to perforated diverticulitis. *Br J Surg* 2008; **95:** 97–101.
12 Sanderson ER. Henri Hartmann and the Hartmann operation. *Arch Surg* 1980; **115:** 792–3.
13 Walklett CL, Yeomans NP. A retrospective case note review of laparoscopic versus open reversal of Hartmann's procedure. *Ann R Coll Surg Engl* 2014; **96:** 539–42.
14 Constantinides VA, Tekkis PP, Athanasiou T *et al.* Primary resection with anastomosis vs. Hartmann's procedure in nonelective surgery for acute colonic diverticulitis: a systematic review. *Dis Colon Rectum* 2006; **49:** 966–81.
15 Zorcolo L, Covotta L, Carlomagno N, Bartolo DC. Safety of primary anastomosis in emergency colo-rectal surgery. *Colorectal Dis* 2003; **5:** 262–9.
16 Gachabayov M, Oberkofler CE, Tuech JJ, Hahnloser D, Bergamaschi R. Resection with primary anastomosis vs nonrestorative resection for perforated diverticulitis with peritonitis: a systematic review and meta-analysis. *Colorectal Dis* 2018; [EPub ahead of print].
17 Schilling MK, Maurer CA, Kollmar O, Buchler MW. Primary vs. secondary anastomosis after sigmoid colon resection for perforated diverticulitis (Hinchey Stage III and IV): a prospective outcome and cost analysis. *Dis Colon Rectum* 2001; **44:** 699–703; discussion 703–5.
18 Fleming FJ, Gillen P. Reversal of Hartmann's procedure following acute diverticulitis: is timing everything? *Int J Colorectal Dis* 2009; **24:** 1219–25.
19 Hallam S, Mothe BS, Tirumulaju R. Hartmann's procedure, reversal and rate of stoma-free survival. *Ann R Coll Surg Engl* 2018; **100:** 301–7.
20 Roig JV, Cantos M, Balciscueta Z *et al.* Hartmann's operation: how often is it reversed and at what cost? A multicentre study. *Colorectal Dis* 2011; **13:** e396–402.
21 Garber A, Hyman N, Osler T. Complications of Hartmann takedown in a decade of preferred primary anastomosis. *Am J Surg* 2014; **207:** 60–4.
22 Feingold D, Steele SR, Lee S *et al.* Practice parameters for the treatment of sigmoid diverticulitis. *Dis Colon Rectum* 2014; **57:** 284–94.
23 Roig JV, Sanchez-Guillen L, Garcia-Armengol JJ. Acute diverticulitis and surgical treatment. *Minerva Chir* 2018; **73:** 163–78.
24 Chow A, Tilney HS, Paraskeva P, Jeyarajah S, Zacharakis E, Purkayastha S. The morbidity surrounding reversal of defunctioning ileostomies: a systematic review of 48 studies including 6,107 cases. *Int J Colorectal Dis* 2009; **24:** 711–23.
25 Garcia-Botello SA, Garcia-Armengol J, Garcia-Granero E *et al.* A prospective audit of the complications of loop ileostomy construction and takedown. *Dig Surg* 2004; **21:** 440–6.
26 Wong NY, Eu KW. A defunctioning ileostomy does not prevent clinical anastomotic leak after a low anterior resection: a prospective, comparative study. *Dis Colon Rectum* 2005; **48:** 2076–9.
27 Kong AP, Kim J, Holt A *et al.* Selective treatment of rectal cancer with single-stage coloanal or ultralow colorectal anastomosis does not adversely affect morbidity and mortality. *Int J Colorectal Dis* 2007; **22:** 897–901.
28 Chun LJ, Haigh PI, Tam MS, Abbas MA. Defunctioning loop ileostomy for pelvic anastomoses: predictors of morbidity and nonclosure. *Dis Colon Rectum* 2012; **55:** 167–74.
29 Biondo S, Kreisler E, Millan M *et al.* Impact of surgical specialization on emergency colorectal surgery outcomes. *Arch Surg* 2010; **145:** 79–86.
30 Patel S, Senagore A. General surgeons vs. colorectal surgeons: who should be doing what to whom? *Semin Colon Rectal Surg* 2013; **24:** 191–4.
31 Moug SJ, Henderson N, Tiernan J *et al.* The colorectal surgeon's personality may influence the rectal anastomotic decision. *Colorectal Dis* 2018; [Epub ahead of print]. https://doi.org/10.1111/codi.14293
32 Wacha H, Linder MM, Feldman U, Wesch G, Gundlach E, Steifensand RA. Mannheim peritonitis index – prediction of risk of death from peritonitis: construction of a statistical and validation of an empirically based index. *Theoretical Surg* 1987; **1:** 169–77.
33 Clavien PA, Vetter D, Staiger RD *et al.* The Comprehensive Complication Index (CCI®): added value and clinical perspectives 3 years "Down the Line". *Ann Surg* 2017; **265:** 1045–50.

Authorship list

Writing group

Luis Sánchez-Guillén (Chair), Dmitri Nepogodiev, Emre Sivrikoz, Tim van Elst, Ana María Minaya-Bravo, Francesco Pata, James Glasbey, Thomas Pinkney, Aneel Bhangu

ESCP cohort studies and audits committee

Alaa El-Hussuna (2017 Audit Lead), Aneel Bhangu, Nicholas Buchs, Christianne Buskens, Sanjay Chaudri, Matteo Frasson, Gaetano Gallo, James Glasbey, Ana María Minaya-Bravo, Dion Morton, Ionut Negoi, Dmitri Nepogodiev, Francesco Pata, Tomas Poskus, Luis Sánchez-Guillén, Baljit Singh, Oded Zmora, Thomas Pinkney (Chair)

Statistical analysis and data management

James Glasbey, Dmitri Nepogodiev, Rita Perry, Laura Magill, Aneel Bhangu (Guarantor)

ESCP research committee

Dion Morton (Chair), Donato Altomare, Willem Bemelman, Steven Brown, Christianne Buskens, Quentin Denost, Charles Knowles, Søren Laurberg, Jérémie Lefèvre, Gabriela Möeslein, Thomas Pinkney, Carolynne Vaizey, Oded Zmora

Collaborators

See page 27–32 for full print list. All collaborating authors will be listed in the online edition, and be PubMed citable.

Correction added on 30 October 2018, after first online and print publication: the author's name Ana Minaya was changed to its full name Ana María Minaya-Bravo.